


UC-NRLF




B 4 175 686

BRARY
IVERSITY OF
ALIFORNIA

TH
ENCES
RARY



EX LIBRIS







THE UNIVERSITY OF CHICAGO PRESS

THE UNIVERSITY OF CHICAGO PRESS

Comparative Heights of Mountains, see p. 105.

Chimborazo

Tunguragua

Mont Rosa

Peak of Teneriffe
Mont Perdu

Lomnits Peak
Kriwan

Elie Mountains

Plomb de Cantal

Ben Nevis

Vesuvius
Snowdon
Helvellyn
Ben Lomond

Level of the Sea



Cotopaxi

Mont Blanc
Jauricocha Silver Mine

Schrekhorn

Mont Tiliis

Etna

City of Quito
Canigon
Kesmark Peak

City of Mexico

Lake of Lançon

Puy de Lanff

Puy de Dome

Schiffellen
Skidaw
Saddelback
Ingleboro

Pyramid of Egypt

Scale 4000 feet to an inch.

W. B.

AN

OUTLINE

OF

MINERALOGY AND GEOLOGY,

INTENDED FOR THE USE OF THOSE

WHO MAY DESIRE TO BECOME ACQUAINTED WITH

THE

ELEMENTS OF THOSE SCIENCES;

ESPECIALLY OF

YOUNG PERSONS.

ILLUSTRATED BY FOUR PLATES.

BY WILLIAM PHILLIPS,

MEMBER OF THE GEOLOGICAL SOCIETY.

NEW-YORK:

PRINTED AND SOLD BY COLLINS AND CO.

.....

1816.

W. Brownell M. D.

E365

P54

EARTH
SCIENCES
LIBRARY

THE UNIVERSITY OF CHICAGO
LITHO

THE UNIVERSITY OF CHICAGO
LITHO

THE UNIVERSITY OF CHICAGO
LITHO

THE UNIVERSITY OF CHICAGO
LITHO

PREFACE.

THE motive for introducing this little volume to public notice, might seem to be wanting if it were not avowed. It is this: There is no elementary work on the subjects it embraces in our language; no book that is calculated, by its simplicity and freedom from theory, and from the shackles imposed upon a learner by the unnecessary use of scientific terms, to invite his attention to the sciences of Mineralogy and Geology.

It is not pretended that the following pages have any claim to originality: all the merit that belongs to them, if indeed any should be allowed, is that of combining in a narrow compass, an outline of sister sciences which merit a more general attention than is given to them; in an arrangement as simple as the subjects will readily allow, and in language which, it is hoped, will be intelligible to those who may have no acquaintance with them.

The form in which this outline is given, that of a division into Lectures, though not absolutely

novel, is not common. During the last winter these Lectures were delivered at the neighbouring village of Tottenham, in the order in which they are now printed; but with some deficiencies supplied, and some errors corrected, that were incidental to hasty compilation. These Lectures were given gratuitously; and the interest they seemed to excite, in a numerous audience, principally composed of young persons, and of both sexes, was felt as a flattering compensation.

But the form of Lectures is not adhered to on that account alone. It allows of a familiarity not inconsistent with an elementary treatise, while it affords an opportunity for useful recapitulation; that perhaps would appear objectionable in any other form; and for occasional repetition, which, if the scientific should condescend to peruse it, may seem tiresome, but will, I have no doubt, be advantageous to the learner.

In a work, by far the greater part of which is compilation, it may reasonably be expected that authorities should be quoted. That has not always been done in its pages. I therefore here acknowledge my obligation, in respect to the mineralogical part, to Aikin's Dictionary of Chemistry and Mineralogy; and in the geological part, to the Transactions of the Geological Society, to Cuvier's Theory of the Earth, edited by Professor Jameson, to whose Geognosy I am scarcely less indebted. These works are my principal authorities; many others were occasionally consult-

ed, but were not made use of in a degree that seems to render their enumeration requisite.

If the perusal of this little volume should tend to create in any person the desire for a knowledge of the sciences on which it treats, beyond their mere elements, it must be owned that it is difficult to refer the reader to works in the English language that are adapted to the use of the learner. The only means by which a knowledge of mineralogy can be acquired, is an acquaintance with minerals: and I have no hesitation in recommending those who may feel this laudable anxiety for further information, in respect to these interesting pursuits, to the acquirement of it by means of small collections, which may be had of one hundred varieties and upwards, with an arranged catalogue, of Mawe, 149, in the Strand, at any price between £5 and £100. One of these little collections would materially assist the progress of the learner; more especially if accompanied by a studious attention to the Manual of Mineralogy, by Arthur Aikin, Secretary to the Geological Society. The introduction to that work, forms a valuable compendium of mineralogical information: and though the learner will meet with many terms and names that will demand explanation, it may be said to be the only work in our language that will be found of advantage to him. The study of geology should follow that of mineralogy. Small collections of rocks may also be obtained.

The Geognosy of Jameson is altogether a scientific work, not well adapted to the learner; inasmuch as a preponderating anxiety for the support of a favourite theory, has caused the introduction of many terms not hitherto adopted by English mineralogists; but much useful and valuable geological information may be gleaned from it.

W. P.

London, 1815.

CONTENTS—MINERALOGY.

	Page
Preliminary observations	1
Objects of Mineralogy and Geology defined	2
Of ELEMENTARY SUBSTANCES	3
Of Mineral substances; simple and compound	4
Explanations of the terms <i>acid</i> and <i>oxide</i> (note)	—
Of AFFINITY	5
the cause of <i>solidity</i> in mineral bodies	6
the cause of <i>crystallization</i> or regular structure	7
Fracture or cleavage of crystals in particular directions	—
Definition of the term, Primitive crystal {	8
The forms of PRIMITIVE CRYSTALS	—
<i>viz.</i> The Parallelopiped	—
Octohedron	9
Tetrahedron	10
Hexahedral prism	—
Rhomboidal dodecahedron	11
Of the internal structure of crystallized minerals	—
Of measuring the angles of crystals—the Goniometer	13
Advantages of the reflecting goniometer	14
Of the NINE EARTHS	15
Three or four of them are metallic oxides	16
Of Silex	—
Alumine	17
Zircon	18
Glucine	—
Yttria	19
Barytes	—
Strontian	20
Lime	21
Magnesia	23

	Page
The Earths found in very different proportions	23
Of the two fixed ALKALIES	24
Of the Alkaline Earths	25
Of the Vegetable Alkali or Potash	—
Of the Mineral Alkali or Soda	—
The Alkalics have metallic bases, and are metallic oxides	26
Of the TWENTY-SEVEN METALS	29
Only seven metals known to the ancients	—
General characters of the metals	—
Of the nature of metalliferous ores	30
Of the MALLEABLE METALS,	
viz. Platina	31
Gold	32
Silver	34
Mercury or Quicksilver	35
Lead	36
Copper	38
Tin	39
Iron	40
Zinc	42
Palladium	43
Nickel	---
Of the BRITTLE METALS,	
viz. Arsenic	44
Antimony	45
Bismuth	46
Cobalt	---
Manganese	47
Tellurium	48
Titanium	---
Tantalium	49
Molybdena	---
Tungsten	50
Chrome	---
Osmium	51
Iridium	---
Rhodium	---
Uranium	---
Cerium	52

CONTENTS—GEOLOGY.

ix

	Page
Of COMBUSTIBLES	52
Of the two bases of Combustibles---Carbon and Sulphur	53
List of Combustible substances,	
viz. Sulphur	—
The Diamond	—
Mineral or Carbon Charcoal	54
Plumbago or Graphite	—
Mineral Oil (naptha and petroleum)	55
Bitumen Mineral or Pitch (maltha and asphalt)	56
Coal (brown, black, and Cannel)	57
Blend or Kilkenny Coal, or Anthracite	58
Jet, or Pitch Coal	—
Amber	—
Mellite, or Honeystone	59

CONTENTS—GEOLOGY.

Of the objects of Geological inquiry	61
The ancients ignorant of Geology	62
Geology, very modern, as a science	—
Hypotheses of Burnet, Woodward, and others	—
More modern theories of Bertrand, Marschall, and Jameson	64
Of the theories relating to the agency of fire and water	68
Principal tracts of low country	71
Strata of low and level countries are horizontal	72
Low and level countries contain sea-shells	—
Sea-shells found in hills far above the sea	—
And at the foot of great chains of mountains	73
Here, the strata are nearly vertical	—
Many catastrophes by the agency of water	74
Summits of lofty mountains contain no shells	—
And are therefore termed <i>primitive</i>	—
GEOLOGICAL POSITIONS	75
Proofs that low and level countries contain organic remains	78
Strata of the Chalk basin of Paris described	79
Strata of the Isle of Wight Chalk basin described	83
Strata of the London Chalk basin described	84
The order of these strata is never inverted	86

	Page
Proofs that catastrophes by water have been extensive . . .	86
Bones of the rhinoceros, hippopotamus, elephant, abundant	88
Found only in alluvial soil in low countries . . .	—
Never in rocks	—
Account of the mammoth found in Siberia (note)	—
Additional proofs of the extensiveness of these catastrophes	90
Rock of Gibraltar encloses bones of the antelope and mouse	—
Bones of the ox, horse, ass, sheep, land shells, and serpents in others	91
Proofs that organic remains are found in hills . . .	96
Cliffs of the Isle of Sheppey enclose fossil fruits, and remains of animals	—
Beds of oyster shells near Reading, and in France, &c.	—
Remains of fish in Germany	97
Shells on the sides of Mount Ætna	—
Proofs that shells are found at the foot of chains of mountains	98
Proofs that the strata, here, are nearly vertical . . .	100
Rocks succeed each other in regular order . . .	101
Secondary rocks known by the shells they enclose . . .	—
Proofs that the summits of lofty mountains contain no shells	—
Of the summits of high mountains	102
The nature and position of their masses	103
Are in their <i>primitive state</i>	104
COMPARATIVE HEIGHTS OF MOUNTAINS	105
Division of rocks into <i>primitive, transition, flötz, and alluvial</i>	108
Primitive rocks are chemical deposites, contain no or- ganic remains	—
Transition rocks are both chemical and mechanical de- posites, and contain organic remains	—
Difference between a chemical and a mechanical depo- site (note)	—
Flötz rocks chiefly mechanical deposites, and contain organic remains	109
Alluvial rocks are mechanical deposites, and contain organic remains	110
Transition and flötz, are by some termed <i>secondary rocks</i>	—
Werner's classification of rocks	112
List of Primitive rocks	—
List of Secondary, or Transition and Flötz rocks	113

	Page
Werner's list of Alluvial deposits	113
Section of the Brocken mountain described	—
Remarks on the strata of mountains	115
Order in their deposition universally prevalent	118
Of PRIMITIVE ROCKS,	
viz. Granite	119
Gneiss	121
Micaceous schistus	123
Primitive limestone	124
Primitive trap	126
Serpentine	127
Porphyry	128
Sienite	129
Topaz rock	130
Primitive flinty slate	—
Primitive gypsum	—
White stone	131
Other rocks are by some added to those called primitive	—
Of TRANSITION, or the OLDER SECONDARY ROCKS	133
viz. Transition limestone	—
Transition trap	—
Greywacke	—
Transition flinty slate	134
Of FLÖTZ, or the NEWER SECONDARY ROCKS,	
viz. Old red sandstone	—
First flötz limestone	—
First flötz gypsum	—
Second, or variegated sandstone	136
Second flötz gypsum	—
Second flötz, or shell limestone	—
Third sandstone	—
Rock salt formation	—
Chalk formation	—
Flötz trap formation	—
Independent coal formation	—
Newest flötz trap formation	—
Of ALLUVIAL DEPOSITES	—
Of organic remains in secondary rocks and alluvial deposits	138
Of MINERAL VEINS	141

Of SALT DEPOSITES	Page
Of COAL DEPOSITES	153
Of VOLCANOES	160
Of the DELUGE	166
Of the INTERNAL STRUCTURE of the EARTH	175
CONCLUDING OBSERVATIONS	181
	187

PLATES.

Comparative heights of Mountains	to face title
Forms of Primitive Crystals	— p. 8
Section of the Brocken Mountain	— p. 114
Veins in Tin Croft and Pink Mines	— p. 149

LECTURE I.

Preliminary Observations—Objects of Mineralogy and Geology defined—Elementary Substances—Simple and Compound Minerals—Affinity—Crystallization—Structure—Primitive Crystals—Of the Earths—Of the Alkalies.

THE outline of the sciences of Mineralogy and Geology, of which I am now about to endeavour to give some idea, is not intended to involve all the nicer inquiries, connected with the subject, that have been instituted by scientific men. Nor do I propose that this outline shall in any degree be dependent upon, or connected with, the many crude and almost barbarous theories of others, who long amused and even dazzled the world by the splendour of their inventions; which tended to retard, rather than to forward an inquiry into the nature of the globe we inhabit.

The phenomena presented by nature, are worthy of our notice; to these your attention will be principally invited.

Of the nature of the globe we comparatively know but little; our investigations are at the best but superficial. For we know nothing but of what appears *on*, or *above*, or of what is brought to light

by the descent of the miner *beneath*, the general level of its surface ; but the miner rarely descends more than 1500 feet, which is little more than $\frac{1}{30000}$ th part of the diameter of the earth. The globe has often been said to resemble in shape, an orange ; in allusion to that resemblance ; we may therefore say, that we know nothing but of the outer rind.

The greater number of mineral substances are to the generality of mankind only rude masses, divested of instruction, and equally unintelligent and unintelligible ; created only to minister to our necessities. To some, it may be even difficult to imagine how they should become the objects of a distinct science, or that after the miner has brought them to light, the naturalist should find an interest in them previously to their being subjected to the ingenuity of the artist.

The sciences of Mineralogy and Geology are, however, worthy of our attention ; they will be found to perform more than they seem to promise. The more we know of them, the more of order, of design, and of contrivance we shall perceive. The power that created the whole is evident in the smallest component part of the most elevated mountain.

Mineralogy has for its object the study of mineral bodies in *particular* ; their characters, varieties, forms, and combinations.

Geology embraces the study of the earth in *general*, of its plains, hills, and mountains, and of

the relative positions of the masses of which they are composed.

Geology comprises the study of rocks in the mass ; Mineralogy, of the individual portions, or substances which, by entering into combination, form the mass. A knowledge of mineralogy is therefore essential to the geologist, and for this reason we shall begin with Mineralogy.

It was anciently supposed, under the dominion of fancy and metaphysics, that all natural substances were ultimately resolvable into four simple bodies, viz. *air, fire, water, and earth*, which hence were called the four elements. The ancients however confessed that the precise natures of the two first, air and fire, were not known to them. They supposed most liquids to be modifications of the third element, water ; and that the solid parts of the globe were attributable to the last, that is, the element they called earth. Thus, combustible bodies they supposed to contain a combustible earth, and metals a metallic earth. To modern chemistry however we are indebted for a much larger catalogue of simple or elementary substances. It is now considered that there are *9 earths, 2 alkalies, 27 metals, and 2 simple substances*, which may be considered as the bases of those termed *Combustibles* ; and *these elementary substances* in the simple or compound state, according to the present state of our knowledge, *form all the various constituent masses of the globe.*

A *simple mineral substance*,* pure Gold for instance, may be described as an *unorganized body*, presenting an assemblage of lesser portions of the same nature, united by the agency or force of a natural law, to which I shall presently advert.

Compound mineral bodies are naturally found in some instances *simply aggregated*, in others *chemically combined*. When simply aggregated, as for instance, when gold occurs in limestone, their separation may be effected *mechanically*, by pounding and washing: but when *chemically combined*, as when silver occurs united with sulphur, we must depend on the labours of the chemist for their separation.

Some of the earths, the two alkalies, and some of the metals, are naturally found combined with various *acids*.†

* A simple mineral substance may be chemically described as a substance that has neither been decomposed nor formed by art.

† Atmospheric, or common air, is composed of two very different kinds of air, or gas, viz. oxygen gas, and azotic, or nitrogen gas, in the proportion of 22 of the former to 78 of the latter.

Oxygen gas, being that part of common air which is essential to animal life, has been also called *Vital Air*: oxygen gas is composed of oxygen united with caloric, or the matter of heat, whence *oxygen* is termed *the basis of Vital air*.

When a substance is combined with a small proportion of *oxygen*, the compound is called an *oxide*: when with a larger proportion, it is called an *acid*, from the sour taste which most of these compounds possess; and chemists have arranged with the acids some compounds which have not been proved to contain oxygen, because their general resemblance to the acids in other respects is sufficient to warrant the opinion that the similarity holds in this particular also.

Some of the metals are also found in combination with *oxygen* which is the basis of vital air; they are then termed *oxides* of those metals; and in that state (to use a familiar illustration) bear the same affinity to those metals, as rust does to iron; rust being the consequence of the absorption by iron, of oxygen from atmospheric air. This chemical combination with oxygen causes metals to assume appearances quite different to the same metals in the pure state; as for instance, the red oxide of copper, is of a ruby red colour, and frequently almost transparent, and the oxide of tin when pure, is nearly colourless and transparent. To show the liability of mineral bodies to become *compound*, it may be quoted that the ore called white silver, is composed of four metals, silver, lead, antimony, iron; two earths, alumine and silex; and one combustible, viz. sulphur.

Water is however found to enter more or less into the composition of some mineral bodies, and it is termed the *water of crystallization* when these substances possess a *regular structure*.

The law to which an allusion was just now made, as being that to which are subject the elementary substances, of which the masses constituting the globe are composed, is called *Affinity*.

When Oxygen is combined with iron, the compound is termed oxide of iron: Oxygen by combining with a certain proportion of carbon or charcoal, forms an acid, called the carbonic acid: which, united with lime, forms a compound mineral called carbonate of lime, of which one variety is limestone, and another is chalk.

This law may be said to involve in it, all those termed attraction, gravitation, magnetism, and electricity. To the law of affinity, mineral bodies owe their existence in separate and similar masses ; as well as their regular crystalline forms ; and but for this important law of nature, the solid parts of the globe would only have been a chaotic mass, instead of exhibiting, even in the oldest rocks, deposits of distinct substances frequently in regular crystalline forms.

Before we proceed to consider the natures of the several simple bodies just alluded to, either separately, or as entering prominently into combination in certain mineral substances, I shall invite your attention to that consequence of the law of affinity, which is termed *Crystallization*. It seems necessary here to introduce this part of our inquiry, because in speaking of earthy and metalliferous substances, I shall present to your notice specimens, some of which exhibit those bodies in their natural crystalline forms. This subject, when considered at large, is curious and interesting ; its investigation, keeping pace with other rapid advances in science, has of late been dignified by the adaptation of geometry and algebra to its illustration. In the narrow compass of what I have undertaken, it is impossible to give more than a very limited view of the subject, but I shall endeavour to make its general outline understood, avoiding, on this, as on all other occa-

sions, as much as possible, all technical or scientific terms.

The term *Crystal*, is derived from the Greek *Κρυσταλλος* (*Crustallos*), signifying ice, which was so called on account of the ease with which it liquified. The term crystal was afterwards applied to what is now called Rock crystal, by the Roman naturalists, and supposing it, both from its transparency and beautiful symmetrical forms, to be only water indurated by continued frosts in the mountainous regions of the Alps. But finding that certain salts also took a prismatic form, the term *crystal* assumed a more general meaning, and now includes all the *regular, many-sided solids*, whether of earthy or metalliferous substances, presented either by nature or chemistry.

There are comparatively very few substances that are not found naturally crystallized with more or less regularity. The forms assumed by some are very numerous and highly interesting. For instance, the carbonate of lime has been found in about 300, and the oxide of tin in about 180 varieties of form: and as of both these, as well as most other substances, the crystals *may be broken in particular directions*, the true definition of a crystal seems to be this, that it has not only *a regular external figure*, but also *a regular internal structure*. This structure in the crystals of several minerals, is readily exemplified by mechanical means, producing fractures, or cleavages of perfect regularity. Some, even small


crystals, have been found exhibiting upwards of 150 little planes; but as most crystals can be broken in particular directions, we are enabled to trace these complicated forms through their many intermediate varieties, into *one simple form*, which therefore is termed the *primitive crystal* of that substance. The crystals of many minerals assume the same figures, and therefore are readily traced to the *same* primitive form. If we take into consideration the whole range of substances found in a crystallized state, the *primitive forms* of their crystals, numerous, varied, and complicated as many of them are, may be said to be comprehended in the five following solids; see plate 2.

The Parallelopiped,
 Octohedron,
 Tetrahedron,
 Hexahedral prism,
 Rhomboidal dodecahedron.

The *Parallelopiped* may be said to comprehend *every solid contained under six parallelograms*: that is, under six planes, whose bounding lines two and two are parallel with each other, and whose opposite sides are equal and parallel. Of these solids, the cube and the rhomboid are selected as the most common: there are also others more or less allied to these. The cube is a body of perfect proportion; all its sides are equal—all its planes are square. The primitive forms of some substances are much flatter than the cube; of others, they are much higher: these are then

Parallelepiped

Cube




Rhomboid




Octohedron

Regular



Acute & Obtuse



Tetrahedron



Hexahedral prism



Rhomboidal Dodecahedron



Faint, illegible text, likely bleed-through from the reverse side of the page. The text is arranged in approximately 20 horizontal lines.



termed *square prisms*. The rhomboid differs from the cube essentially; none of its planes are square, but they resemble each other; two of the opposed angles of each plane are more obtuse, the other two are more acute, than those of the planes of the cube. Of rhomboids there are several varieties; that is, the angles of their planes are more or less acute and obtuse: these rhomboids therefore differ in a greater or less degree from the proportions of the cube. There are some other varieties of the parallelepiped which it does not seem essential to our present object to explain.

Of the *Octohedron* also there are several varieties: three of the most obvious are selected. The *regular octohedron*, is so called, because it is a solid contained under eight planes, each of which is a triangle equal and similar to each of the others; and which, having all its three sides of the same length, is therefore termed an *equilateral triangle*. A line drawn from its summit to its lowest extremity or angle, is precisely of the same length as a line from any other of its angles to its opposite angle. As the primitive crystal of some substances, is found an octohedron, much flatter than the regular, and therefore called an *obtuse octohedron*: two of the three lines, bounding each of its eight triangles, are shorter than the third. A line drawn from the summit to its lower extremity or angle is shorter than a line from any of its other angles to its opposed angle. An octohedron is found to be the

primitive crystal of some substances which is much longer than the regular octohedron, and is called *acute*, because its upper and lower angles are more sharp or acute than the other four angles. Two of the three lines bounding each of its eight triangles, are longer than the third; and a line drawn from the upper to the lower extremity or angle, is much longer than one drawn from either of its other angles, to the opposed angle.

Of the octohedron there are some other varieties, of which it does not seem now requisite to enter into an explanation.

The *Tetrahedron* is a solid comprehended within four planes, each of which is a triangle equal and similar to the rest, having each of its three sides of the same length. It is the simplest of all the primitive forms.

The *Hexahedral prism* is a solid comprehended within eight planes, which are not all similar. Its upper and lower planes, called the *terminal* planes, are alike, being six-sided or hexagonal: each of the six sides being of the same length. The other six planes, which are called the *lateral* planes, are alike in regard to each other, and in the figure in plate 2, are each perfectly square. This is not always the case in primitive crystals; in some, the lateral planes are much longer, in others shorter; in either of which cases the planes would not be square, but oblong. Hexahedral prisms, as primitive crystals, are therefore of various lengths.

The *Rhomboidal dodecahedron* is a solid comprehended within twelve planes, all of which are perfectly alike in form; each plane being a rhomb, of which the four bounding lines are all equal in length, and are parallel two and two, forming two acute and two obtuse angles on each plane.

But, certain substances are naturally found to assume two of those, which are termed *primitive forms*. Amongst these is the fluete of lime; it therefore becomes a question which of the two, is the *true* primitive form of that substance. This question is decided by the *internal structure* of the crystal; if we take a crystal of it in the form of the *cube*, we shall find that all its corners, termed the solid angles, may be regularly broken off, so as to produce in lieu of each of them, a regular triangular plane, and that, by pursuing this fracture, we shall finally arrive at a form which in fact is the *regular octohedron*, which therefore is proved to be the *primitive* form of the fluete of lime; of which, it is also proved, that the *cube* is only a *secondary* crystal, the result of an aggregation of laminae on the several planes of the octohedron; in other words, by the addition of laminae, or regular layers, parallel with the planes that may be obtained by mechanical cleavage.

If however we take a similar crystal, that is, a cube, of common salt, we shall find that it is not the result of the same law of crystallization; that is, we shall find that it cannot be fractured in the same way, the corners or angles cannot be taken

off, and as it can be fractured only in the *direction of, or parallel with the six faces* of the cube; that solid is therefore esteemed to be the primitive form of common or rock salt. But by adding regular layers parallel with the planes obtainable by mechanical cleavage, that is, on the six planes of the cube of rock salt, we should never obtain an octohedron; accordingly rock salt is never found crystallized in the form of the octohedron.

The carbonate of lime is always readily broken into *Rhomboids*, and no other form.

The crystals of the fluuate and the carbonate of lime, and of common salt, are quoted as remarkable instances of the ease with which their natural structure affords mechanical cleavage; but, all mineral substances are not fractured with the same ease. The topaz can only be readily cleaved in one direction; and quartz or rock crystal, though sometimes it presents indications of its real structure, has never yet been regularly fractured. This diversity in the structure of crystallized bodies has not been satisfactorily explained; we only know the fact.

It has already been said that the mineralogist depends on the internal structure, for a knowledge of primitive forms; but where the knowledge of this structure cannot be attained by mechanical means, recourse must be had to other means: analogy seems the only resource. In other words, a comparison of the external forms of such crystals as do not admit of a mechanical cleavage, with

the external forms of the crystals that do admit of it; but this is sometimes the source of error.

It may readily be supposed that the regular internal structure so observable in the crystals of many substances, would necessarily produce a regular external form. This actually exists in a degree scarcely credible by those who have not examined the subject, both in those crystals of which the internal structure is known, as well as in those of which it is not.

Much attention has lately been given to ascertain precisely, the angles at which the various and even numerous planes of crystals meet each other. For this purpose an instrument has been invented, called the *goniometer*, meaning the measurer of angles; but we may conclude that its use scarcely admits of perfect accuracy; and for this reason, the few most conversant in its use, have differed in the results obtained from the same substances.

Another instrument has lately been invented by Dr. Wollaston, called the *reflecting goniometer*, because its use depends on the reflections to be observed upon the natural polish on the planes of crystals. This instrument has already done service to science in detecting some fallacies, arising from a reliance on the former instrument.

When once the angles of the primitive crystal of any substance are accurately ascertained, the angle at which any two of its numerous planes meet, is calculated with wonderful precision by the assistance of geometry.

The angles formed by the meeting of any two planes of each of the five regular solids, the cube, the regular octohedron, the tetrahedron, the hexahedral prism, and the rhomboidal dodecahedron, already enumerated among primitive crystals, are known, because they all are *regular geometrical solids*. But it has already been shown, that there exist as primitive crystals, *varieties of the parallelepiped and of the octohedron*, which differ from each other both in shape and measurement. As instances; the primitive crystal of quartz, is a rhomboid very different to that of carbonate of lime, and much more nearly approaches the cube. The primitive crystals of the oxyd of tin and of zircon, are octohedrons, differing from each other, but are much more flat than the regular octohedron, while the octohedron which is the primitive crystal of sulphur, is much longer and more acute. The primitive crystal of the cyanite is a four-sided and oblique prism, with rhombic terminations, while that of the sulphate of barytes is a four-sided rhomboidal prism also with rhombic terminations. The difficulty therefore is, to obtain the precise admeasurements of those primitive crystals, which are *not regular geometrical solids*. It is probable that the reflecting goniometer of Dr. Wollaston will discover, that the principal part of the calculations hitberto made, in regard to the angles of such primitive crystals, and of the numerous facets to be observed on them, (which not belonging to the primitive crystals, are

therefore termed *secondary planes*,) are incorrect; and for this reason; this goniometer will assist in measuring the angles of crystals as small as the head of the smallest pin, if their planes be perfectly brilliant; and it is ascertained that minute crystals are always more perfectly formed than large ones. Now the calculations already alluded to are grounded upon measurements taken by the common goniometer upon large crystals, the planes of which rarely are perfectly formed and smooth: even such as have brilliant planes, and seem therefore adapted to the use of the reflecting goniometer, seldom afford similar results.

I shall now offer to your notice some observations on those elementary substances termed simple or *pure earths*, which are nine in number; they are called

- | | |
|------------|--------------|
| 1. Silex | 6. Barytes |
| 2. Alumine | 7. Strontian |
| 3. Zircon | 8. Lime |
| 4. Glucine | 9. Magnesia |
| 5. Yttria | |

All these *earths* agree, when free of foreign admixtures, in this one character, they are all *snow-white*.

Although these substances by common consent are called simple or pure earths, it seems essential, in speaking of them as elementary substances, to say that chemistry, which, during the last twenty years has made amazing progress, has by the late brilliant discoveries of Sir Humphrey Davy,

completely turned the tables on the ancients. They supposed that every metal had an *earthy basis*, whereas his experiments have decisively shown, that three if not four of the earths have a *metallic basis*; that, in fact, they are *not simple substances*, but *compounds*, consisting of metallic bases united with oxygen; and, to use the same familiar illustration as before, these four earths have the same affinity to their respective metallic bases as rust has to iron; *they are metallic oxides*. The four earths are those called silex, alumine, lime and barytes, but the basis of alumine is not so decidedly ascertained as the bases of the other three. Strictly therefore we ought to diminish the number of earths, by the three or four in question, and to increase the number of metals; but, as discoveries, however brilliant and however well established, are rarely admitted with instantaneous consent, these *metallic oxides* are still suffered to hold their places as *earths*.

I now proceed to give some account of each earth separately, and shall offer to your inspection some specimens of substances in which they prominently enter into combination.

The first on the list is **SILEX**. This is one of those earths which the discoveries of Sir H. Davy have decidedly shown to have a metallic basis, which he has called *Silicium*, and being the result of a combination of oxygen with that metal, is therefore a metallic oxide. Silex, in its pure state, is not three times as heavy as water, and has neither taste nor smell.

As common flints are almost wholly composed of *siliceous earth*, it has from that circumstance received the name of *Silex*, which in Latin signifies *Flint*; but this earth is found in greater purity in opal, and in quartz, or rock crystal.* *Silex* is probably the earth which most abundantly enters into the composition of the globe.

Silex enters largely as a component part of glass, for which purpose the pure sand from some parts of the coast of Norfolk, and Alum Bay in the Isle of Wight, are preferred. It has never been found combined with an acid.

ALUMINE has been so called from its forming the basis of common alum. When pure, it has neither taste nor smell, and is twice as heavy as water. Though one of the most extensively diffused substances, it is nowhere found pure; and though a constituent part of the purest clays, it rarely forms more than one-fourth or one-third part of them; it is however found nearly pure in the oriental ruby and the sapphire, and in corundum, which are the next in hardness to the diamond.† Alumine forms a large proportion of that valuable mineral called fullers' earth, which has that smell when breathed on, peculiar to

* Quartz is composed of silex, with 2 or 3 parts in the 100 of water. Opal, of 90 parts of silex and 10 of water. Flint, of 97 parts of silex, 1 of alumine and oxide of iron, and 2 of water.

† By analysis, the *Sapphire* yields about 93 parts of alumine, 1 of oxide of iron, and a small portion of lime; the *Oriental Ruby*, 90 parts of alumine, 7 silex, and some oxide of iron: *Corundum*, about 89 parts of alumine, 5 of silex, and nearly 2 of oxide of iron.

clayey substances, and which forms a mineralogical test of the presence of alumine.

In useful purposes, alumine enters largely into the composition of bricks, pottery, and porcelain; it is infusible. The experiments of Sir H. Davy have rendered it probable, that alumine is a metallic oxide, having a metallic basis, which he has called *aluminum*.

ZIRCON when pure, is rough to the touch, insipid, and insoluble in water; it is found combined with other substances in the hyacinth, from a brook called Expailly, in Auvergne in France; and in the jargoon from Ceylon.* Hitherto zircon has not been put to any useful purpose; it is about four times as heavy as water.

The earth called GLUCINE obtained that name from the Greek *γλυκος*, signifying *sweet*, on account of the sweet taste by which its salts are distinguished. When pure, glucine is a white powder, soft, and somewhat unctuous to the touch, and is nearly three times the weight of water.

In the natural state glucine has hitherto only been found in the beryl and the emerald;† but neither the pure earth nor any of its salts, have hitherto been applied to any use.

* The *Hyacinth* is composed of 70 parts of zircon, 25 of silex, and a trace of oxide of iron. *Jargoon* 66 parts of zircon, 31 of silex, and 2 of oxide of iron.

† The *Emerald* is composed of about 64 parts of silex, 16 of alumine, 13 of glucine, 1 of lime, and nearly 4 of oxide of chrome.

YTRIA, in many of its properties and appearances in its pure state, bears considerable affinity to glucine, it has the same saccharine taste, but is easily distinguished from it, inasmuch as it is nearly five times heavier than water, and by some properties discoverable only by the chemist.

In the natural state, yttria occurs as a component part of a mineral substance called the gadolinite,* which is brought only from Sweden, and which was so called on account of its having been first analysed by the Swedish professor Gadolin, who named the earth yttria, because the mineral in which it was discovered, was brought from Ytterby in Sweden.

BARYTES has never been found pure in the natural state, but always combined, either with the *sulphuric acid* forming sulphate of barytes or heavy-spar; or with the *carbonic acid* forming carbonate of barytes, or witherite.† Barytes and all its salts, except one, are violent and certain poisons, destroying animals by inflaming the intestines, and are often used for the destruction of vermin. Barytes is one of the earths that have lately been discovered by Sir H. Davy to have a metallic basis. This basis, he has called *Barium*,

* The *Gadolinite* is composed of about 56 parts of yttria, with a trace of manganese, 22 of silice, 5 of glucine, 17 of oxide of iron, a small portion of alumine and some water.

† The *Witherite* is composed of 78 parts of barytes and 22 of carbonic acid. *Heavy Spar* is composed of 87 parts of barytes and 33 of sulphuric acid.

which uniting with a certain proportion of *oxygen*, forms barytes.

STRONTIAN has never been found in a pure state. It was first brought, combined with the *carbonic acid*, from a place in Argyleshire called Strontian, whence the mineral obtained its familiar name of strontianite. Strontian, combined with the *sulphuric acid*, is found near Bristol; thus combined, the mineral substance has obtained the name of cœlestine,* from its delicate tint of a light blue colour.

The earth called strontian, either when pure or combined with the acids above mentioned, has not hitherto been applied to any use. At present therefore these combinations merely serve to form a part of a mineralogical collection.

LIME has never yet been found in a pure state, but when so prepared by the chemist, is moderately hard, and of a hot acrid taste. It has been proved by Sir H. Davy to have a metallic basis which he calls *Calcium*: lime is therefore a metallic oxide. Lime naturally occurs in combination with the *carbonic, sulphuric, boracic, fluoric*, and *phosphoric acids*.

Lime, combined with the *carbonic acid* forms a mineral substance, thence termed carbonate of lime.† A variety of this mineral is to be found in

* The *Strontianite* is composed of about 69 parts of strontian, 30 of carbonic acid and a little water. *Cælestine*, of 54 parts of strontian, and 46 of sulphuric acid.

† *Calcareous Spar*, or carbonate of lime, is composed of 57 parts of lime, and 43 of carbonic acid.

the cabinets of mineralogists, in a vast number of beautiful forms, all of which may be traced to the rhomboid, which therefore is termed the primitive crystal. Those minerals commonly called limestones, and chalk, and marble, are also carbonates of lime. The uses to which lime is applied, (when the carbonic acid with which it is combined, as forming chalk and limestone, is driven off by heat) are well known. The rocks on each side the Avon near Bristol, are of a peculiar kind of limestone, which has obtained the name of swine-stone, from its yielding when rubbed a fetid smell. This smell is attributed to the presence of bitumen, the nature of which will be mentioned among the combustibles.

Lime, in combination with the *sulphuric acid*, is called *sulphate of lime*, or familiarly *gypsum*.* Of this mineral the uses are very extensive. When compact, it is called *alabaster*, and is employed by the architect for columns and other ornaments, being more easily worked than marble; it is also turned by the lathe into cups, basins, vases, and other similar articles. When sulphate of lime or gypsum, is subjected to a certain heat, it loses what is termed its water of crystallization, and is converted into a fine powder called *plaster of paris*; the uses of which, when beaten up with water into a paste, for taking casts of gems and statues, are well known.

* *Gypsum* is composed of about 32 parts of lime, 46 of sulphuric acid, and 21 of water.

Lime, combined with the *arsenic acid*, forms a mineral called *Pharmacolite* ;* the only species : with the *boracic acid*, a mineral is formed called the *datholite*.†

Lime, combined with the *fluoric acid*, is called *fluuate of lime*. This mineral is commonly known by the name of *fluor spar* ;‡ in Derbyshire, as *blue john*. The elegant vases and chimney ornaments turned by the lathe out of blocks of this substance, are known to almost every one ; the brown colour generally observable in these vases is obtained by exposure to heat. Fluuate of lime is extensively used in smelting the ores of copper.

The mineralogist values the fluuate of lime for its abundant and beautiful variety of crystals, both in form and colour : all the forms it assumes may be traced to the regular octohedron, which therefore is termed the primitive form of its crystal. The octohedron may be obtained from each of them, however unlike to it, by breaking it in certain directions.

A variety of fluuate of lime, called *chlorophane*, found in Siberia and Cornwall, on being exposed to the heat of a live coal, gives out a beautiful green light.

* The *Pharmacolite* is composed of about 46 parts of arsenic acid, 23 of lime, 6 of silex, 22 of water, and a small portion of oxide of cobalt.

† The *Datholite* is composed of about 36 parts of lime, 36 of silex, 24 of boracic acid, and 4 of water.

‡ *Fluor Spar*, or the fluuate of lime, yields by analysis about 68 parts of lime and 32 of fluoric acid.

Lime, in combination with the *phosphoric acid*, forms a mineral which occurs in small crystals, called *apatite** or phosphate of lime.

MAGNESIA is a light earth of a perfect whiteness, and is absolutely insipid; the slightly acrid taste occasionally to be found in the magnesia used in medicine, arises from a proportion of lime. Naturally, magnesia occurs combined with the *boracic acid*, in the borate of magnesia, or boracite.†

In the rare mineral by some called native magnesia, analysis has proved it to be combined with the sulphuric and carbonic acids.

Magnesia is one of the least abundant substances; it does not enter into the composition of many mineral bodies, but it forms a small proportion of that substance called the *soap-stone*, which owes its greasy or soapy feel to magnesia. The soap-stone is largely employed in the manufactory of porcelain.

These nine earths enter, in very different proportions, into the composition of the globe.

It is considered that silex is the most abundant of all. It forms the greatest ingredient of the oldest rocks, is largely found in others, and in

* *Apatite* yields 55 of lime and 45 of phosphoric acid.

† The *Boracite* is composed of about 17 parts of magnesia and 83 of the boracic acid. The *Spinelle Ruby*, about 85 parts of aluminic, 9 of magnesia, 6 of chromic acid. *Asbestus*, 59 parts of silex, 25 of magnesia, 9 of lime, 3 of aluminic. The *Soap-stone*, of about 59 parts of silex, 30 of magnesia, 2 oxide of iron, and 6 of water.

clays and soils; in these, alumine is the next in abundance: to it succeeds lime, which is less common in primitive rocks, though very plentiful in the transition and flötz, or secondary rocks.

Magnesia and barytes occur in comparatively very small quantities. The first enters but little into the composition of rocks and soils; the latter rarely.

Strontian, zircon, glucine, and yttria, are very sparingly found; the first may be said to be the most common of the four, the others are only found in part, the components of a few mineral substances, some of which are occasionally enclosed in rocks; but rarely does any one of these four earths enter into the composition of rocks or of soils.

Barytes, magnesia, strontian, and lime, are never found pure; but mostly combined with acids.

We now proceed to those mineral substances termed ALKALIES, which enter into the composition of several minerals.

The term *Alkali* is Arabic, and is expressive of the acrid saline residue left in the ashes of the plant called *Kali*, after its combustion in the open air; and, not being volatile, was termed *Fixed Alkali*.

FIXED ALKALIES are usually denominated of two kinds. The *Vegetable* or Potash, and the *Mineral* or Soda. *Potash* is procured from the ashes of vegetables in general *not* growing contiguous to the sea. *Soda* is the basis of common

salt, and is therefore found in immense quantities; it is also the principal saline residue of plants growing *on* the sea shore.

The taste of the pure fixed alkalies, is acrid, burning, and nauseous; they are without smell; they have peculiar chemical properties, which it is not my province to describe; but it is important to say that the earths called barytes, strontian, lime, and magnesia, agree so nearly with the alkalies, in some of their chemical properties, that some chemists have given them a place among the alkalies, others have termed them alkaline earths.

POTASH occurs in the natural state in the leucite, and in mica, and in several other substances; therefore the term vegetable, as applied to potash, is not absolutely correct.*

SODA is found naturally combined with the carbonic acid, forming carbonate of soda, and with the muriatic acid, forming muriate of soda or common salt.†

The mineral substance called fettstein, yields, by analysis, both potash and soda.‡

Both potash and soda are largely employed in the making of glass and soaps. When pure, they

* The *Leucite* yields, by analysis, about 54 parts of silex, 25 of alumine, and 21 of potash.

† The water of the ocean contains from one-twenty-fifth to one-thirty-fifth of its weight of muriate of soda or common salt, which is composed of 53 parts of soda, 47 of the muriatic acid.

‡ The *Fettstein*, 44 silex, 34 alumine, 4 oxide of iron, a small portion of lime, and about 16 parts of soda and potash.

are not easily distinguished from each other, but potash is the heaviest. Until lately, both potash and soda were believed to be simple substances; but, by the experiments conducted by Sir H. Davy, with the astonishing powers of electro-chemical agency, they have both, as well as the four earths already noticed, been actually decomposed. The mode by which this was effected is highly ingenious and interesting; the detail belongs to the chemist. The result, however, was, that *potash* was found to be a combination of *oxygen* with a *metal*, exactly resembling quicksilver in appearance, and which, by the discoverer, has been called *Potassium*, as being the basis of potash. The result was exactly similar in regard to *soda*, the metallic basis of which he called *Sodium*. Both potash and soda, therefore, are metallic oxides. Both of the metals, *Sodium* and *Potassium*, are malleable; in this respect, agreeing in character with iron, copper, and some other metals.

The same observation that has been made in regard to the four earths, which have metallic bases, viz. that their bases properly belong to the metals, will also apply to potassium and sodium, but potash and soda still hold their rank as fixed alkalies.

It is by no means improbable, that future researches in chemistry, will also discover the bases of the remaining five substances yet considered as earths, and that, by common consent, both the

earths and the fixed alkalies will be swept away, and their bases added to the catalogue of metals.

The discoveries just noticed have been effected by the aid of galvanism superadded to chemistry. Galvanism, or electricity, therefore, (for a doubt of their identity scarcely exists) holds a high and important station in science. With such aid, chemical experiment, conducted by such men as Sir H. Davy, and other able chemists of the present day, may work a complete reformation in science: it is impossible even to conjecture where discovery will stop.

LECTURE II.

Of the Metals—Of Combustibles.

ON a former evening it was said that mineral bodies are considered to be of four kinds, **EARTHS**, **ALKALIES**, **METALS**, and **COMBUSTIBLES**.

That they are naturally found either simple or compound; the simple consisting of one substance alone, the compound of two or more, and that these are either mechanically or chemically combined.

That some of the earths and metals are found combined with various acids, and some of the metals with oxygen, or the basis of vital air.

It was also said that, by the aid of chemistry, nine earths had been discovered, and two fixed alkalies; but that Sir Humphrey Davy, by the assistance of electro-chemical agency, had lately proved that three or four of those substances called pure earths, as well as both the fixed alkalies, have, in fact, metallic bases; and that they consist of oxygen in combination with those bases.

Having also, in the former evening, considered some of the properties and uses of the earths and alkalies, we shall now proceed to notice the metallic bodies, and afterwards those called combustibles.

The only metals known to the ancients were gold, silver, copper, iron, tin, lead, and mercury; but discoveries have from time to time increased the catalogue, until it has been swelled to the number of twenty-seven, independently of those which have very lately been discovered as the bases of four of the earths and the two alkalies.

Of these metals, the first eleven only have the important property of malleability, or of being sufficiently tenacious to bear the extension of their body by beating with the hammer; the others have by some been therefore termed brittle metals.

Malleable Metals.

Platina,
Gold,
Silver,
Mercury,
Lead,
Copper,
Tin,
Iron,
Zinc,
Palladium,
Nickel.

Brittle Metals.

Arsenic,
Antimony,
Bismuth,
Cobalt,
Manganese,
Tellurium,
Titanium,
Tantalium,
Molybdena,
Tungsten,
Chrome,
Osmium,
Iridium,
Rhodium,
Uranium,
Cerium.

A lustre is peculiar to the metals, which, therefore, is called the metallic lustre: another remarkable property is their want of transparency when in the mass; but, as leaf gold, held between the eye and a luminous body, transmits a green light, and silver a white light, it seems probable that other

metals, if attenuated in the same degree, would also be translucent.

In weight the metals far exceed the earths; the heaviest of the earths is only about five times heavier than water, but the lightest of the metals is more than six times heavier than water. Beaten gold is nineteen times heavier than water; and beaten platina, the heaviest of all, is twenty-three times heavier than water.

The characters of fusibility and extensibility in metals, is of vast importance to man; for without these characters neither could they be freed from the earths and other impurities with which they are naturally found; nor without these characters could they be wrought into vessels for his use.

Metals are believed to be *simple substances*; *not one of them has hitherto been decomposed.*

The only metals that, as yet, have been found in the pure or *native* state, are platina, gold, silver, quicksilver, copper, antimony, arsenic, tellurium, bismuth, and iron. But the greater part of these are rarely found quite pure, but mostly have small proportions of other metals intermixed.

In order to illustrate the very brief view I am about to take of the several metals, I shall offer to your inspection specimens of some of them as they are naturally found either in their simple state, (if so found) or as forming those compounds which are denominated *metalliferous ores*.

An *ore* is a compound of two or more metals, or of a metal in combination with oxygen; (whence

such a combination has obtained the name of a *metallic oxide*;) or a metal combined with an *acid*, or with a *combustible*. Many ores are of so compound a nature as to consist of two or three metals united with oxygen, sulphur, one or more of the earths, and with water.

I now proceed to invite your attention to the metals individually; beginning with those which possess the qualities of fusibility, ductility, and malleability, so important to man.

PLATINA is about twenty-three times heavier than water; hitherto it has only been found in Peru, Brazil, Spain, and in the island of St. Domingo. In Peru, it is found in little flattened grains, rarely exceeding the size of a pea, accompanied by gold, and the ores of titanium and iron; yet it is said that Humboldt presented the King of Prussia with a mass larger than a pigeon's egg. But the grains of *crude platina* are not pure; analysis has proved them to consist of platina alloyed by four other metals, osmium, iridium, rhodium, and palladium, which will hereafter be noticed. The platina of Brazil is alloyed by gold and silver; that of Spain was found in a gray silver ore.

Crude platina is very difficult of fusion. Pure platina, in thin plates, is very ductile and flexible. Of late it has been formed into mirrors for reflecting telescopes, spoons, crucibles, and some vessels of considerable dimension for the use of the che-

mist in particular processes. So ductile is platina, that Dr. Wollaston has lately succeeded in drawing it into a wire $\frac{1}{18736}$ part of an inch in diameter.

GOLD, when pure and beaten, is about nineteen times heavier than water, is very soft, and perfectly ductile and flexible. So great is the tenacity of gold that a piece one-tenth of an inch in diameter, will hold five hundred pounds without breaking; and it is computed that a single grain of gold will cover the space of fifty-six square inches, when beaten out to its greatest extent.

Gold is mostly found in the metallic form, whence, by mineralogists, it is said to occur in the native or pure state; but it is generally alloyed by small portions of other metals, as silver, copper, &c. It occurs in mineral veins and beds, or disseminated in the substance of some of the oldest mountains: it is found in Brazil, Peru, Mexico, Africa, Sumatra, Japan, Hungary, and Transylvania.

Helms says, that when a projecting part of one of the highest mountains in Paraguay fell down, about thirty years ago, pieces of gold, weighing from two to fifty pounds each, were found in it; and that in the Vice-royalty of La Plata alone, there are thirty gold mines.

Sometimes gold is crystallized in small cubes or regular octohedrons, and as these crystals cannot be broken in any particular direction, either of

those solids may be said to be the primitive crystal of gold. In veins it is generally accompanied by quartz, felspar, the ores of tin, silver, lead, and of some other metals.

A great quantity of gold is obtained in grains and rounded masses in soils, evidently the ruin of rocks which contained it in its natural situation. In this state it has been found in Wicklow, in Ireland, and in Cornwall, in small quantities. A few years ago, a single specimen of gold, equal in weight to upwards of ten guineas, was found among tin in a stream work in Cornwall. On the coast of California there is a plain of fourteen leagues in extent, about fourteen inches beneath the surface of which, large lumps of gold are irregularly interspersed.

But a still greater quantity of gold has been obtained in the form of a fine sand, from the Peruvian, Mexican, and Brazilian rivers, and from some of the African. In Europe, the Danube, the Rhine, and the Rhone, and the streams of Hungary and Transylvania, afford small quantities.

The uses of gold are well known. Alloyed by copper, it is employed for ornamental purposes, coin, and plate.

In English coin it was alloyed by two parts of copper to twenty-two of gold. The purple colour used in porcelain painting is obtained from a preparation of gold.

SILVER, when pure, is ten times heavier than water, and is soft, opaque, and flexible; a piece one tenth of an inch in diameter will support two hundred and seventy pounds without breaking.

Silver naturally occurs in the pure or native state; but is sometimes alloyed by a small proportion of gold, sometimes of copper. It is found in fine filaments disseminated through rocks, but chiefly in veins, in primitive and secondary mountains, occasionally crystallized in cubes and regular octohedrons, and accompanied by calcareous and other spars, iron pyrites, cobalt, and some other substances. It is found in Peru, Mexico, Saxony, Bohemia, Norway, Hungary, and England.

The ores of silver are numerous; for although it mostly occurs in the pure state, it is also found combined with gold, copper, antimony, iron, lead, bismuth, arsenic; with the earths, silex, and alumine, and mineralized by the *muriatic*, *sulphuric* or *carbonic acids*, and by *sulphur*.

The most common ore of lead, called the sulphuret, mostly contains some portion of silver, but not always worth extracting. The lead from the Westmoreland and Cumberland mines, yields an average of seventeen ounces of silver to the ton of lead. The Beeralston mines in Devonshire, yield about forty-two ounces. The richest perhaps ever known, was that found at Brunghill Moor, in Yorkshire; which yielded two hundred and thirty ounces to the ton.

According to Helms, the mine of Jauricocha, in Peru, which is about three miles above the sea, contains a prodigious mass of porous brown iron-stone, half a mile long, as much broad, and about one hundred feet in depth, which is throughout interspersed with pure silver, and a white argillaceous vein very much richer. It is asserted that Jauricocha, and the mines of the district surrounding it, have yielded forty millions of dollars in a year.

It is said that, in 1750, a mass of silver was found in a mine near Freyberg, in Saxony, weighing upwards of 140lbs. and another of about the same size, in 1771. In the year 1748, a block of native silver and silver ore was cut out in a rich vein of silver, near Schneeberg; Duke Albert, of Saxony, descended the mine, and used it as a dinner table. When this huge block was smelted, it yielded 44,000lbs. of silver. We are told that Annibal received 300 pounds weight of silver, daily, from the mines near Carthagena, in Spain.

The uses of silver are numerous, and for the most part obvious. In coin, silver is alloyed by one part of copper to fifteen of silver. The yellow colour used in porcelain painting is oxide of silver.

MERCURY or **QUICKSILVER**, is thirteen times heavier than water, and is fluid in the natural temperature of the atmosphere.

It mostly occurs pure (but sometimes contains a little silver) disseminated in globules, or collected in the cavities of its mines, which are most

commonly situated in calcarerous rocks, or indurated clay, or argillaceous schistus.

Quicksilver mines are worked in Carniola, the Duchy of Deux-ponts, Spain, and Peru. The vein of Guancavelica, in South America, in which quicksilver is found in the state of cinnibar, is 80 Spanish ells in extent, and is situated partly in sandstone, partly in limestone. The cinnabar is accompanied by the sulphuret of lead, calcareous spar, barytes, quartz, manganese, arsenic, &c.

The quicksilver mines of Idria, in Saxony, are said to yield 100 tons annually; and those of Spain a still greater quantity.

The ores of mercury are not numerous: combined with silver, it is called *native amalgam*, with sulphur and iron, *cinnibar*. *Horn mercury* is a natural combination of mercury mineralized by the *sulphuric acid*, and of mercury mineralized by the *muriatic acid*.

The uses of mercury in medicine, in the arts, and in experimental philosophy, are numerous; but its chief use is in the separation of gold and silver from their ores, by a process called amalgamation. When amalgamated with tin, and laid on glass, it forms mirrors.

LEAD, when pure, is more than eleven times heavier than water; a piece one-tenth of an inch in diameter, will hold twenty-nine pounds without breaking.

Lead has never yet been found pure in the natural state. Its ores are numerous, and occur

in beds or veins in almost every mineral district in the known world, and are, perhaps, next to those of iron, the most common of metalliferous ores.

Lead is found in combination with other metals, as antimony, iron, and silver; and the two earths, silex and lime. It is found mineralized by the *carbonic, muriatic, phosphoric, arsenic, molybdic, and chronic acids*, and with *oxygen*, which cause it to lose every appearance and character of lead; but many of its ores have not been analysed.

The most common of the ores of lead, is by far of the greatest importance to man, because, from it are principally derived the immense quantities of lead for his use. It is called galena, or sulphuret of lead; by analysis it yields lead, sulphur, oxide of iron, and sometimes lime and silex; mostly some silver. It occurs in beds or veins in primitive or secondary mountains, most abundantly in argillaceous schistus and secondary limestone, accompanied by blende and calamine, the ores of zinc; and is sometimes compact, sometimes crystallized in the cube or regular octohedron.

It would scarcely be possible to enumerate all the valuable purposes to which lead is applied in the arts, in medicine, and in the common wants of man. Among its less obvious uses, lead is employed to glaze pottery, and its oxide enters into the composition of glass. Four parts of lead and one of antimony form printing types; to which, by some, is added a little copper or brass.

With tin and bismuth it forms alloys mentioned in the notice of those metals.

COPPER, in its pure state, is about eight times heavier than water; a wire one-tenth of an inch in diameter will support two hundred and ninety-nine pounds and a half without breaking.

It is a very malleable and ductile metal, of a pale red colour, with a tinge of yellow. In the natural state it occurs very pure; and its ores are very numerous. In both states, it is found in almost every mineral district in the world, in beds, or more commonly in veins, in primitive and secondary mountains, accompanied by several other mineral substances, as the ores of zinc, and occasionally of lead, sometimes of tin; with quartz, and fluate of lime, and calcareous spar in abundance.

Native or pure copper, is not, however, found either in beds or veins in great quantities; that of Japan and of Brazil is alloyed by gold. A mass of native copper is said to have been found in a valley in Brazil weighing 2666 Portuguese pounds. Wherever found, it is of various shapes, and sometimes crystallized in the cube and regular octohedron.

Mineralized by a certain proportion of *oxygen*, it forms a beautiful mineral, called the *red oxide of copper*, which assumes a great variety of forms, all of which may be traced into the regular octohedron; with an increased proportion of oxygen, it assumes a black hue, and is mostly pulverulent.

Copper is found combined with iron and arsenic, with lime and silex, and mineralized by the *phosphoric, carbonic, arsenic, or muriatic acids*, which cause it to lose all metallic character and appearance.

The most common copper ore of the Cornish mines is of a yellow colour, called yellow copper ore, or copper pyrites; analysis proves it to consist of copper, iron, and a large proportion of sulphur.

The uses of copper, in all its various states, are almost endless, and only, if at all, inferior to those of iron. Alloyed with certain proportions of zinc it forms brass, pinchbeck, tinsel, and Dutch gold, in imitation of gold leaf. With a small proportion of tin, copper forms bronze or bell metal; but if the proportion of tin amount to one-third, it forms speculum metal, used for reflecting telescopes. With zinc and iron, it forms tutenag. In porcelain painting, the green is obtained from copper.

TIN, in its pure state, is about seven times heavier than water, but has never been found pure. In the common ore of tin mines, it is always in combination with *oxygen*, whence it is termed an oxide; but analysis proves it also to contain small portions of iron and of silex.

In one vein in Cornwall, an ore has been found called the *bell metal ore*, (from its resemblance to that metal in colour,) which consists of *tin, copper, and sulphur*, together with a small portion of *iron*.

A variety of the oxide of tin, called *wood tin*, is found sparingly in two or three places in Cornwall only.

Tin is considered to be one of the oldest metals, because it is principally found in those rocks which, from their not containing any animal or vegetable remains, are termed primitive. It occurs disseminated in them, or in beds, but principally in veins, accompanied by the ores of tungsten, arsenic, iron, copper, and zinc, and with quartz, mica, fluate of lime, and some other substances.

The ore of tin is also abundantly found in Cornwall, in rounded portions or grains, in what are termed alluvial beds; that is, in depositions which have resulted from the ruin of rocks.

Tin is by no means one of the most commonly diffused metals. It is most abundant in Cornwall; but it is also found in Gallicia, in Spain, in Saxony, in Bohemia, in Malacca and Banca, in Asia, and in Chili, in South America.

The alloys of tin with other metals, are mentioned in treating of lead, copper, and quicksilver. Another will be noticed under the article bismuth. In a fine leaf, as tin foil, it is used for many purposes; its salts are used in dying: its economical purposes are well known.

IRON, when pure, is about seven times heavier than water; it is one of the most, if not the most, universally diffused substances in nature; it is found in all soils, and in almost every rock.

Iron has been said to have been found in a mine in Saxony, in the pure or native state, alloyed by small proportions of lead and copper; but the fact has not been satisfactorily ascertained.

But some masses of a substance, which by some is termed *native iron*, have been found in different quarters of the globe; in Bohemia, in Senegal, in South America, and in Siberia; of the latter we have the best account. It was found by professor Pallas on the top of a mountain, on which there was a considerable bed of magnetic ironstone, on the banks of the river Jenisei. It weighed 1680 Russian pounds, and possessed some of the important characters of pure iron, as malleability and flexibility, and was reported by the inhabitants of the country to have fallen from the sky. The mass found in the Vice-royalty of Peru, in South America, was described by Don Rubin de Celis: it weighed about fifteen tons; it was compact externally, and was marked with impressions as if of hands and feet, but much larger, and of the claws of birds; internally it presented many cavities: it was nearly imbedded in white clay, and the country round it was quite flat and destitute of water.

Most of these masses termed native iron, (which from a current belief of their having fallen from the sky, have also obtained the names of *meteoric iron*) have been subjected to analysis, and in each the iron has been found alloyed with more than one-tenth of the rare metal called *nickel*; which

also, it is worthy of remark, is found, by analysis, to be a constituent part of all those stones, which, in various parts of the European Continent, in England, and in America, have been known to fall from the sky, and are therefore termed *meteoric stones*.

The ores of iron are numerous, and are found in beds, in veins, and disseminated in rocks. It occurs combined with *manganese, carbonate of lime, silex, alumine, sulphur, or oxygen*: with *copper, the arsenic acid, oxygen and silex*, it forms a beautiful mineral crystallized in cubes of a green colour, which are often transparent; it is called the *arsenate of iron*. An ore, in which iron is combined with alumine, is used in the making of what are termed *red lead pencils*. Plumbago, or *black lead*, is a natural compound of iron, with a large proportion of carbon.

It would be vain to attempt the enumeration of the uses to which iron is put by man. Steel is an artificial combination of iron with carbon. The red colour used in porcelain painting is oxide of iron.

ZINC, when pure, is about seven times heavier than water; its tenacity is not great; a piece one-tenth of an inch in diameter will hold twenty-six pounds without breaking; and being far less ductile than some other metals, its importance is thereby diminished.

Zinc is never found pure, its ores are only three in number, but of these some varieties are found

Zinc, as an *oxide*, combined with *carbonic acid*, forms a most abundant ore, called *calamine*. Zinc, as an *oxide*, combined with *silex*, forms *electric calamine*, so termed from its becoming electric when slightly heated. Zinc, combined with *iron*, *sulphur*, *silex*, and *water*, forms that ore called *blende*; a variety of which, on being scratched, emits a phosphoric light.

The ores of zinc are found in most mineral countries; most abundantly in the transition or earlier secondary rocks, accompanied by iron pyrites, sulphuret of lead, some of the ores of silver, and by calcareous spar and quartz.

Zinc is employed by the Chinese for coins: it enters into the composition of many alloys; (see copper.) It is sometimes used in medicine, and in oil painting.

PALLADIUM, is about eleven times heavier than water; it is very malleable, and equal in hardness to bar iron. It has hitherto only been obtained by the chemist from *crude platina* (see Platina,) which it greatly resembles in colour: it has never been applied to any use.

NICKEL is about nine times heavier than water, and is of a yellowish white; it is not perfectly malleable.

Nickel has never been found in the pure state: its ores are few; they have been found in mineral veins and beds in France, Spain, Bohemia, Siberia, and in England sparingly; they are generally accompanied by the ores of silver and cobalt, by

calcareous spar and quartz, and some other substances.

It is remarkable that nickel, which is one of the least abundant metals, has been found, by analysis, to enter into the composition of those stony substances which, in various parts of Europe and America, have fallen from the atmosphere; whence they are termed meteoric stones.

The uses of nickel are not numerous; it is chiefly employed in alloys with other metals.

We have now taken a slight view of the eleven metals, which have been termed *perfect*, on account of their possessing the valuable properties of *fusibility*, *ductility* and *malleability*. I now proceed to those which, not possessing the two latter properties, have been by some termed the *brittle*, or *semi-metals*.

ARSENIC is nearly eight times heavier than water, and is of a bluish white.

It is found nearly pure, being alloyed only by small portions of iron and sometimes of gold or silver, only in primitive mountains, in veins, accompanied by some ores of silver, cobalt, and lead; by calcareous spar, fluate of lime, and quartz, and some other substances. It is principally found in certain districts of Germany.

Arsenic, combined principally with *iron*, forms a mineral called *arsenical pyrites*, or *mispickel*; in some of the varieties of which, silver is found.

This also principally occurs in veins in primitive mountains.

Arsenic is found combined with twenty-five parts of *sulphur*, forming an ore of a red or orange colour, called *realgar*, and with forty-three parts of *sulphur*, it forms an ore of a bright lemon yellow colour, called *orpiment*. *Realgar* is said to occur principally in primitive mountains; *orpiment* principally in flötz or secondary mountains.

Arsenic is one of the least useful metals, and though a poison, is used in medicine; it is also used in the making of glass: *orpiment* is employed as a paint.

ANTIMONY is a compact, brittle, whitish metal, about six times heavier than water. It is found nearly pure, being alloyed only with very small portions of silver and iron. Native or pure antimony is found in veins in the mountains of Dauphiné, in the Hartz, &c. and in Sweden, disseminated in calcareous spar.

The ores of antimony are only five in number; all of which have not been analysed. In some of them, it is found combined with oxide of *iron*, *cobalt*, *arsenic*, *silex*, *sulphur*, *oxygen*. They are found principally in veins in primitive, and in transition, or the older secondary, mountains in Sweden, Saxony, France, Bohemia, England, and other mineral countries.

Antimony forms alloys with other metals, and is used in the arts. It enters largely into the com-

position of printing types ; it is also used in medicine.

BISMUTH is nearly 10 times heavier than water ; it is of a reddish-white colour, and very brittle.

It is found in the pure state somewhat alloyed by arsenic.

The ores of bismuth are only two in number ; in that called *sulphuret of bismuth*, it is combined with *sulphur* ; in the other, called *bismuth ochre*, it is mineralized by *oxygen*, and combined with small portions of *oxide of iron* and *carbonic acid*.

Native bismuth is rare, as well as its ores ; these are found in veins mostly in primitive mountains, accompanied by the ores of cobalt, of iron, of zinc, and sometimes of silver, and by quartz, calcareous spar, and barytes ; in Bohemia, Transylvania, France, and Sweden. The sulphuret of bismuth has occurred in Cornwall.

Bismuth is very little used, but it enters into the composition of some of the soft solders, and of sympathetic ink. It forms alloys with other metals. Tin and bismuth are two of the most fusible metals. The fusible-metal of Sir Isaac Newton, is composed of 8 parts of bismuth, 5 of lead, and 3 of tin ; when this is thrown into water, and heat applied, it melts a little before the water has reached the boiling point.

COBALT, when pure, is about 8 times heavier than water : it is of a gray colour, with a red tinge, and has the magnetic properties of iron.

It is not found pure. Its ores are not numerous. In one of them from Tunaberg, in Sweden, it is combined with *arsenic* and *sulphur*, and somewhat in form and colour, resembles iron pyrites. In Cornwall, it is found combined with *arsenic* and *iron*. Its other ores have not been analyzed, but cobalt seems always to be combined with arsenic.

The ores of cobalt occur in veins both in primitive and in secondary mountains: mostly accompanied by some of the numerous ores of copper, sometimes by native bismuth, native silver, native arsenic, and the ores of silver.

Cobalt is very little used except in the arts. It is brought to this country reduced to the state of an oxide, of an intense blue colour, called *zaffre*, which, when melted with 3 parts of sand and 1 of potash, forms a blue glass, and when pounded very fine, is called *smalts*, and is then employed to give a blue tint to writing papers, and in the preparation of cloths, laces, linens, muslins, &c.; for colouring glass, and for painting blues on porcelain. So intense is the blue of *zaffre*, that one grain will give a full blue to 240 grains of glass.

MANGANESE is of an iron gray colour, very brittle, and seven times as heavy as water.

It is never found pure. Its ores are not numerous. It is found combined with the *oxide of iron*, with *sulphur*, the *sulphuric* or *carbonic acid*, *barytes*, or most abundantly with *oxygen*, as an oxide of a brown colour or black.

The ores of manganese are found in various parts of the continent of Europe, and in the mineral districts of Britain.

From the black oxide of manganese, all the oxygen gas used by the chemist is obtained, and all the oxygen entering into the composition of the oxymuriatic acid consumed in the bleacheries of Britain, France, and Germany. The violet colour employed in porcelain painting is obtained from manganese. In glass-making, it is employed in the finer kinds of glass, both as a colouring material and a destroyer of colour: this application of it is ancient; it is mentioned by Pliny.

TELLURIUM, when pure, is about the colour of tin. It is brittle, nearly as fusible as lead, and is six times heavier than water.

It is an extremely rare metal, and is found in the native state, but always alloyed by other metals, principally in veins traversing secondary rocks in Transylvania. A variety called the *gray gold ore* of Nagyag, has been found, by analysis, to consist of *tellurium, lead, gold, silver, copper, and sulphur*.

It has never been made any use of.

TITANIUM is so difficult of fusion, that the attempts to reduce it to a pure metallic state, have scarcely succeeded.* It is of a copper red colour.

Two of its ores are said to be nearly pure *oxides*. In others, is found in combination with *oxide of iron, manganese, and silex*. They occur sparingly

in Hungary, Transylvania, France, Britain, and North and South America.

The hair-like appearances sometimes to be observed in crystals of quartz, are mostly crystals of titanium. An ore of titanium is found in a stream in Cornwall in black grains; another is found in Transylvania resembling yellow sand.

The only use to which titanium has ever been put, was in the porcelain manufactory at Sevres, where it was employed to produce the rich browns in painting it. The want of uniformity in colour occasioned its disuse.

TANTALIUM is a metal, having but a slight external metallic lustre; it is dull, and almost black internally.

It is extremely rare; having hitherto only been found in Finland and Sweden. In Finland, it occurs combined with *oxide of iron* and of *manganese*, forming a mineral called *tantalite*, imbedded in quartz, in veins that traverse a red granular felspar. In Sweden, it is found in a mineral called *yttrotantalite*, because analysis has proved it to be principally composed of the rare earth *yttria*, and the rare metal *tantalium*. This mineral occurred in a granite rock.

MOLYBDENA, when pure, is of a grayish white, and in the form of brittle infusible grains.

It is very rare, and has never been found pure. Combined with *sulphur*, it is found in veins in primitive mountains, in Norway, Sweden, Saxony,

and Switzerland, accompanied by tin, wolfram, quartz, and mica.

The *molybdic acid* has been found combined with *lead*, forming a mineral called *molybdate of lead*, in Carinthia, Saxony, Hungary, and Austria, accompanied by calcareous spar, sulphuret of lead, the ores of zinc, and fluor spar.

Molybdena has never been applied to any use.

TUNGSTEN is a hard, brittle, granular metal, of a light steel-gray colour, and brilliant metallic lustre.

It is not found pure.

The *oxide of tungsten*, combined with *lime* and *silex*, has been sparingly found in Sweden, Bohemia, and Cornwall. The compound is called *tungstate of lime*.

Combined with *oxide of iron*, *manganese*, and *silex*, it forms a mineral called *tungstate of iron*, or *wolfram*, which occurs in most districts in which tin is found.

The only use to which tungsten has hitherto been applied is in the arts, as forming, in combination with other substances, those red paints known by the name of lake.

CHROME, is a metal of a grayish-white colour, and extremely brittle; it is remarkable that it has never been found in the *metallic form*, but only in the *acid state*, or in that of an *oxide*.

The *chromic acid*, in combination with *lead*, forming a compound mineral called *chromate of lead*, has been found principally, in veins in gneiss

and mica-slate, in a gold mine in the Uralian mountains, in Siberia : it is said also to have occurred at Annaberg, in Austria, and at Trapettes, in Savoy. It is extremely rare.

The *oxide of chrome*, in combination with *oxide of iron*, *alumine* and *silex*, forms a mineral called *chromate of iron*, which is found very plentifully in France, and in some places in Siberia.

The chromate of lead, on account of its beautiful red colour, has been employed in Russia as a paint. Chrome, as obtained in the metallic state by the chemist from either of the two foregoing compounds, has not been applied to any important use : it tinges glass of a green colour. It has been ascertained that the emerald owes its beautiful green colour to *oxide of chrome* : it seems, therefore, probable, that chrome may hereafter be employed as a paint.

OSMIUM,

IRIDIUM, and

RHODIUM, are three brittle metals which, together with PALLADIUM, already noticed as a malleable metal, have, by analysis, been found in combination with PLATINA in the crude state, (see Platina.) Not one of these four metals has hitherto been applied to any use.

URANIUM, is a brittle, granular, hard metal, of extremely difficult fusibility.

It is remarkable that this metal has never been found in any state having a *metallic* appearance ; consequently, never in the pure state. Its ores

are only two ; and very rare. They have been found in Saxony, Bohemia, Norway, France, and England ; in the latter only in copper veins.

Uranium is found only in combination with oxygen, forming an *oxide* ; or with oxygen and oxide of iron, when it is called *uran-ochre*.

The oxide of uranium is a beautiful mineral, mostly in small thin plates of a fine green colour, and transparent : in combination with oxygen and oxide of iron, uranium forms a mineral similar in appearance to pitch ; or sometimes resembling iron-rust.

No use has hitherto been made of uranium.

CERIUM has hitherto only been obtained by some chemists from a mineral substance from Sweden, called the *cerite*, in the form of a white, yellowish, or reddish brown powder : yet, although cerium has never been obtained in the metallic form, it is considered from its properties to be a metal.

It has not been applied to any use.

We now proceed to the consideration of those mineral bodies which, from their peculiar properties, are termed **COMBUSTIBLES**. These form, in the mineral kingdom, a class of substances by no means agreeing amongst themselves in internal or external characters, and differing essentially from the earths, the alkalies, or the metals. Combustibles include both the hardest and the softest of mineral substances.

Most of the metals, whose properties are altered by combustion, acquire an *increase* of weight thereby; whereas combustible substances are sensibly *diminished* in weight by the same process. The product of some of them is liquid, of others, solid; if solid, it is insoluble in water.

The *mineral bases* of combustible substances may be said to be only two, viz. CARBON and SULPHUR.

Combustible substances may be comprised in the following list :

Sulphur	Coal
Diamond	Blind or Kilkenny Coal,
Mineral Carbon	or Anthracite
Plumbago, or Graphite	Jet
Mineral Oil	[Pitch Amber
Bitumen, or Mineral Mellite,	or Honey Stone.

SULPHUR, is a soft, brittle substance, of a pale yellowish colour. It is found either nearly pure, or in combination with certain metallic ores, in great abundance. It is also found both in the vegetable and animal kingdoms.

The DIAMOND, which is the hardest substance in nature, was heretofore considered as an earthy or stony substance; but it is proved beyond a doubt not to be an earthy substance. When exposed to a current of air, and heated to the temperature of melting copper, it is found to be gradually, but *completely* combustible. By this pro-

cess, it may be wholly converted into carbonic acid, and therefore consists of *pure carbon*.

Diamonds are either colourless, or of a yellowish, bluish, yellowish green, clove brown, black brown, prussian blue, or rose red colour:—they are naturally found in detached regular crystals; their primitive form is the regular octohedron. In India, diamonds are found in an indurated ochrey gravel. The diamond mines extend through a long tract of country, from Bengal to Cape Comorin: the chief of them are now between Golconda and Masulipatam. Diamonds are also procured from the isle of Borneo and from Brazil, where they are found in beds of ferruginous sand.

The principal use of the diamond is in ornamental jewellery; it is also employed by glaziers to cut glass, and by lapidaries to engrave the harder gems.

MINERAL CARBON, or CHARCOAL, is grayish black. It occurs in plates or irregular pieces, in various sorts of common coal. It has a glimmering, silky lustre, and a fibrous appearance, discovering a wood-like texture. It is somewhat heavier than common charcoal, and is easily reduced to ashes before the blowpipe, without either flame or smoke.

PLUMBAGO, or GRAPHITE, is found in England, Scotland, France, Spain, Germany, and some other countries. Plumbago is of a dark iron black, passing into steel gray; it has a glistening metallic lustre.

The principal use of plumbago is in the making of what are called *black-lead* pencils; for which purpose none has yet been discovered equal to that from Borrodale, in Cumberland.

Whence this mineral obtained the name of *black-lead* it is difficult to say, unless it was from the lead-coloured streak which it gives upon paper. It has been ascertained that lead does not enter into its composition, but that the purest plumbago consists of about 90 parts of carbon and 10 of iron.

MINERAL OIL. Under this term are comprehended two substances, *naphtha* and *petroleum*; both of which are liquid, highly inflammable, and lighter than water.

Naphtha is nearly colourless and transparent; it gives out much smoke and a penetrating odour in burning. The most copious springs of *naphtha* are on the coast of the Caspian sea. It is employed externally for strains and bruises. The Persians and Russians are said to take it as a cordial.

Petroleum, at the usual temperature, is rather thicker than common tar, and has a strong, disagreeable odour. It is principally found in coal countries, as in Colebrookdale. It is found in France, Italy, Switzerland, Germany, Hungary, and Sweden.

It is most plentifully found in Asia: round the town of Rainanghong, in the Birman empire, there are 520 wells in full activity, into which petroleum flows from over coal. No water ever

penetrates into these wells. The quantity of petroleum annually produced by them amounts to more than 400,000 hogsheads. To the inhabitants, its uses are important; it serves instead of oil for lamps, and, mixed with earth or ashes, for fuel.

BITUMEN, or MINERAL PITCH, is either *elastic* or *compact*.

Elastic bitumen, is of various shades of brown. It has a slightly bituminous odour, and is about the weight of water. It burns readily with a large flame and much smoke, but melts by a gentle heat, and is thereby converted into a substance resembling petroleum.

Hitherto it has only been found in the Odin mine, near Castleton, in Derbyshire, in a secondary limestone.

Compact bitumen is of a brownish-black colour; one variety may be impressed by the nail, and is called maltha; another is very brittle, and is called asphalt.

They consist of carbon, earth, and bitumen.

The softer variety has not been put to any use; but the harder is used in varnishes, and is an essential part of those used by engravers. It is found on the shores of the Dead Sea, in the West Indies, and many other places.

COAL. The bituminous substance called coal, though ranked among minerals because its basis is pure carbon, is now, by many, believed to be of vegetable origin; because the substance

which lies upon the coal, is always filled with vegetable remains; as well as because a woodlike appearance may be traced through every species of coal, even the most compact.

On the subject of coal deposits, particularly our own, it is my intention to treat more at large.

Coal may be divided into three species: *brown coal*, *black coal*, and *cannel coal*.

Brown coal is in. perfectly bituminous; in all its varieties it is fibrous, and in some of them its vegetable origin is so complete, as that the remains of the trunks and branches of trees are visible, and almost perfect. Brown coal burns with a weak flame and disagreeable odour. It is found in horizontal strata. In England it occurs at Bovey, near Exeter, and is called Bovey coal. It is also found in other countries.

Black coal, which is used for economical purposes, includes several varieties. It may, however, generally be said to be of a black colour, having an iridescent tarnish, and a high resinous lustre. It is composed of about 60 parts of carbon, and 4 of bitumen. It always occurs in nearly horizontal strata, which are abundant in Durham; Lancashire, Yorkshire, and in some other parts of England, and in several parts of Europe.

Cannel coal is chiefly found at Wigan, in Lancashire, but is more or less abundant in most collieries. It is very brittle, of a shining lustre, it crackles and flies while burning, flames much,

and burns quickly, leaving only 3 or 4 parts in the 100 of ashes.

BLIND, or KILKENNY COAL, or ANTHRACITE, is of a dark iron black, and has a bright, metallic lustre. It burns without smoke, and emits no sulphureous or bituminous odour. It consists of pure carbon, with some silex, and a small portion of oxide of iron.

JET, or PITCH COAL, is generally of a velvet black; it occurs in mass, and sometimes in the shapes of branches, with a regular woody structure. It has a brilliant, resinous lustre. It is used as fuel, but the finer and harder pieces are worked into trinkets, under the name of jet. It is found in the Prussian amber mines in detached fragments, and is there called black amber.

AMBER is a mineral of a yellow, or reddish-brown, or of a greenish, or yellowish white colour. It is found in nodules, or rounded masses, from the size of coarse sand to that of a man's head.

Ambre is found on the shores of the Baltic, of Sicily, and of the Adriatic Sea, and occasionally in the gravel beds near London. Near the sea-coast, in Prussia, there are regular mines of amber: under a stratum of sand and clay, about 20 feet thick, succeeds, a stratum of trees, 40 or 50 feet thick, half decomposed, impregnated with pyrites and bitumen, and of a blackish-brown colour. Parts of these trees are impregnated with amber, which sometimes is found in stalactites depending from them. Under the stratum of trees was

found pyrites, sulphate of iron, and coarse sand, in which were rounded masses of amber. The mine is worked to the depth of 100 feet, and from the circumstances in which the amber is found, it seems plain that it originates from vegetable juices. Amber sometimes incloses insects, believe'd to be of the ant species. The strong electric powers of amber are generally known.

Amber yields, by distillation, an acid called the *succinic acid*, and leaves, as the residue, an extremely black, shining coal, which is employed as the basis of the finest black varnishes. When exposed to flame in the open air, amber takes fire, and burns with a yellowish flame, giving out a dense, pungent, aromatic smoke, and leaving a light, shining, black coal.

The MELLITE or HONEYSTONE is a rare mineral, having hitherto only been found in Thuringia, in the district of Saal, and in Switzerland. It occurs on bituminous wood, and earthy coal, and is generally accompanied by sulphur.

The honeystone is softer than amber, is transparent, brittle, and electric, and is found crystallized in the octohedron.

When burnt in the open air, neither smoke nor flame are observable, and it eventually acquires the colour and consistence of chalk. By analysis, it gives a peculiar acid, called the *mellitic acid*, in combination with alumine, together with small portions of iron and silex.

We have now concluded the subject. What has been said was intended only to convey the elements of Mineralogy. There exists a vast multitude of mineral compounds, which it was not possible to offer to your notice in so short a space of time, and which involve many inquiries and researches belonging more properly perhaps to the mineralogist in his closet.

It may be observed, that, although we should now have been ignorant of the existence of some of the earths and metals, but for the researches of the chemist, yet, the properties and uses of by far the greater number of those which are most useful to man, were known, and employed in his service, long before chemistry was pursued as a science. Perhaps not one half of the earths, metals, and combustibles, have, in any important degree, been hitherto turned to advantage.

LECTURE III.

Of the objects of Geological inquiry—Hypotheses—Geological positions
—Of the low and level parts of the Earth—Of the chalk basins of
Paris, of London, and of the Isle of Wight.

It was said on a former evening that the object of mineralogy is the study of mineral bodies in *particular*, whether simple or compound: that geology embraces the study of the globe *in general*, and of the various relations that the different masses of which it is constituted, bear to each other. Mineralogy may be, therefore, said to furnish, as it were, the alphabet to geology.

The globe we inhabit is about 8000 miles in diameter, 25,000 in circumference. Its surface has two grand divisions, land and water: one-third, or thereabouts, being occupied by land, and two-thirds by water.

So little was known by the ancients respecting the earth, (of its real form and unceasing revolutions they were absolutely ignorant.) that it was by them considered to be the centre of the universe; of which, a more correct philosophy has proved it to be only a subsidiary portion. It is not, therefore, to be wondered at, that in the pursuit of our present object, we derive little or no benefit from the writings of ancient profane

authors. In the time of Herodotus, the Greek historian, it may, however, be inferred that there existed some philosophers who imagined that the earth was round, and that it was encompassed by the sea, since the historian takes the opportunity of sneering at the opinion.

Geology, or the study of the earth, may be said to be altogether modern, as a science. Until towards the end of the last century, it was little understood; perhaps, because those sciences on which it greatly depends, chemistry and mineralogy, had not made any large advances towards their present state. It is no marvel, therefore, that in default of a knowledge of the sciences, and of that research by which alone we can become acquainted with the constituent masses of the globe, the activity of the human mind should attempt to account for the creation and present state of the earth by uninstructed efforts of the imagination. It may be amusing to give a short account of a few of these, broached by men calling themselves philosophers.

In these hypotheses, two events only, the creation and the deluge, seem to have entered into the calculations of the inventors; as comprehending all the changes to which the globe has been subjected: that is to say, each arbitrarily ascribed to it a certain primitive state, which each supposed to be altered and modified by the effects of the deluge.

In the opinion of *Burnet*, the whole earth at

first consisted of an uniform light crust, which covered the abyss of the sea ; and which, being broken for the production of the deluge, formed the mountains by its fragments.

According to *Woodward*, the deluge was occasioned by a momentary suspension of cohesion among the particles of mineral bodies ; the whole mass of the globe was dissolved, and the soft paste became penetrated by shells.

Whiston fancied that the earth was created from the atmosphere of one comet, and deluged by the tail of another.

The great *Leibnitz*, as well as *Descartes*, amused his imagination by conceiving the world to be an extinguished sun or a vitrified globe ; upon which the vapours, condensing in proportion as it cooled, formed seas, and afterwards deposited calcareous strata.

Demaillet conceived the globe to have been covered with water for many thousand years ; that it gradually retired ; that all the terrestrial animals were originally inhabitants of the sea ; that man himself began his career as a fish.

Buffon imagined, that the mass of our earth, together with those of the other planets, were struck off the sun, in a liquified state, by a comet, at the same instant.

Some modern philosophers have supposed, every thing to have been originally fluid ; that this universal fluid gave existence to animals of the simplest kind ; that in process of time the

races of these animals became complicated, and dying, supplied calcareous earth or lime; that aluminous earth or clay, was supplied by the decay of vegetables. That these two earths were re-dissolved by a final analysis into silex; hence that the more ancient mountains are silicious. Thus the solid parts of our globe, according to these visionaries, owe their existence to animal or vegetable life; and without it, would have continued entirely liquid.

Kepler, one of the greatest of astronomers, considered the globe to be possessed of living faculties and a circulating vital fluid; that all the particles of it are alive, and possess instinct and volition, whence their attraction and repulsion: that the organs through which the huge animal breathes are the mountains; that mineral veins are abscesses, and metals the products of rottenness and disease.

These systems, and even many more than these, have had their admirers, and have successively sunk into disrepute and neglect in proportion to the advance of chemical and physical science. It is the apology, if indeed it be not rather the shame of their inventors, that they knew nothing of mineralogy, or of the structure of the earth. Two or three theories, of a date much later than the foregoing—of the end of the last and beginning of the present century, are worth notice. *Marschall* supposes the fragments of which the surface of the earth is composed to have fallen from heaven.

Bertrand has supposed that the earth is hollow, and contains within it a load-stone, which is dragged from one pole to the other by the attraction of comets; so as, by changing its centre of gravity, to drown alternately the two hemispheres. Jameson, now a professor of natural history in one of our own universities, has lately published this amusing query: 'As the true figure of the earth is still unascertained, may we not conjecture, from what is already known, that it is a polyedron (a figure of many sides,) and that the strata, under determinate angles, form the sides and cleavage of this great crystal?'

If philosophers, and even naturalists, will still condescend to amuse the world with conceits like the foregoing, it is no wonder that the present period, with respect to the theory of the earth, should have been said to bear some resemblance to that in which some philosophers thought that the heavens were formed of polished stone, and that the moon was no larger than Peloponnesus. Almost equally absurd, they merit equal notice, as examples of extravagant theory; but they convey to us at the same time this instructive lesson, that it is *only by the patient investigation of facts and of natural phenomena*, that we can hope to approach the truth, in the sublime study of the history of the planet we inhabit.

Within the last few years mineralogy has made advances so rapid, as well as other investigations

connected with the history of the earth, that geology is thereby greatly raised in rank among the sciences. A review of modern researches and discoveries will strikingly evince the folly of any attempt to account, by a turn of the pen, for the creation of the globe, or for the revolutions to which it has been subjected: these researches amongst the phenomena of the earth, will also assure us, that, in whatever manner the mighty display of Omnipotence in its creation was effected, it has since suffered great changes on its surface.

The elementary substances which have been developed by modern chemistry, as entering in a greater or lesser degree into the composition of mineral bodies, consist of nine earths, twenty-seven metals, two fixed alkalies, and the two bases of combustible substances, carbon and sulphur.

It is probable that chemistry will yet make great alterations in the catalogue of elementary substances, either by addition or diminution; most probable by the latter. We shall, however, assume these, according to the present state of our knowledge, to be *the elements of which the various constituent masses of the crust of our globe are compounded*; but, whether the globe is to its very centre a mass of these compounds, we know not: and since there seems no probability of our arriving at any certainty on this head, let us forbear to conjecture. It is the business of the geologist to investigate *natural facts and phenomena*; and let us be assured that when these fail

to accomplish any desirable purpose, we shall add nothing to our knowledge by soaring into the regions of fancy. On a former evening it was said, that as the miner rarely descends below $\frac{1}{1000}$ th part of the earth's diameter, so, if we suppose the earth in shape to resemble an orange, it may be said that we know nothing but of the outer rind.

Geology, therefore, in the present true sense of the term, embraces little more than an *inquiry into the history and present state of the surface or crust of the globe*. Disclaiming all theory, it is my purpose to adhere to the legitimate objects of science, as they have been just described, without exerting one pretension beyond them.

Much has been written on the subject of the creation; a subject so far beyond our limit, that upon it I shall be silent; nor shall I attempt to follow Kirwan through a laborious undertaking to reconcile our *partial knowledge* of the phenomena of nature, and of the constituent masses of the globe, with the Mosaic account of the creation. If ever it should be permitted that man shall comprehend the great plan by which the Creator reduced to order the materials of which the globe is composed, with the same certainty as he has attained a knowledge of the mechanism of the universe, some more certain data and discoveries will be allowed to him, than as yet have fallen to his lot. Geology is yet in its infancy: shall we not, therefore, do wisely in concluding that we are

not in the possession of materials sufficient for the investigation? We have no reason to disbelieve that this important branch of geological inquiry will hereafter be better understood. Considering the present state of our knowledge, we shall be best and most reasonably employed in the investigation of the present state of the earth, and of the changes to which its surface has been subjected; taking as our text, the incontrovertible truth that, 'In the beginning God created the heaven and the earth.'

Some of the most acute geologists of the present day, have not yet agreed, whether the *agent* employed in the magnificent work of reducing to order the heterogeneous mass of elementary substances, was *fire* or *water*. There exist, at this moment, two distinct parties, distinguished according to the notion they embrace, by the appellations of *Volcanists* and *Neptunists*. Each loudly asserts the preference of its own theory; each perhaps in the possession of chemical facts, or of natural phenomena, establishing, in its separate opinion, its own claim to preference, that have not been controverted by the other. Distinct, however, from these two parties, there are many who wisely think that we are yet too ignorant to be able, with propriety, to establish any theory at all. These are collecting evidence in regard to its actual state from every quarter of the globe, which, if faithfully recorded, may

hereafter enable mankind more nearly to approach the truth than at present it is possible.

To the results of these inquiries, in so far as they have proceeded, it is my present object to invite your attention. They will comprehend much on the consequences of the catastrophes that have befallen the surface of the earth by the agency of water; of which we have abundant and incontrovertible evidence: inquiries into the nature and component masses of mountains, and of their relative heights; the internal structure of the earth; mineral veins; the deposits of salt and of coal; and of volcanoes.

These are inquiries well meriting our attention; they will exhibit to us the investigations of men, who, being philosophers in the true sense of the word, have sought and still continue to seek among the instructive records of revolving ages, amid the ruin occasioned by time and circumstance, the history of the globe we inhabit.

He who resides in a country which exhibits an almost perpetual verdure, uninterrupted by barren rocks, or desolate regions, and has never visited any other, will scarcely be led to suspect that the surface of the globe has been much convulsed by successive revolutions and various catastrophes. If, however, he descend to any considerable distance beneath its surface, or ascends the hills that border the plain, he will be likely to receive a new train of ideas; his mind will become

expanded in proportion to the expansion of his view. But if he ascend the elevated chains of mountains, or follow the beds of descending torrents, which lay open their interior structure, he will become prepared to believe, to their full extent, that various catastrophes have befallen the globe since its creation.

We reside in a country just now alluded to ; a country of perpetual verdure, uninterrupted by a single rock, or one desolate place. The elevations that surround us scarcely merit the name of hills ; and being all alike verdant, they admit not of investigation, beyond what is attained in the sinking of wells. Few of us have visited other countries ; not many have seen the more mountainous part of our own ; scarcely one present, perhaps, knows the internal history of the spot which now supports him. We generally know that it principally consists of clay, even to a considerable depth ; but the greater part have yet to learn that this clay has unquestionably been deposited by the sea, that it encloses sea-shells, and that the whole country surrounding us to a considerable depth and extent, is the *debris*, or ruin of rocks.

It seems to me that we cannot do better than begin our inquiries into the nature of such countries as those in which *we* live—of low and level countries ; from which we may rise to the consideration of those somewhat more elevated, and

thence to that of the nature and component masses of mountainous regions.

But level countries are so little open to the investigation of the geologist; and seem to afford, when compared with the more obvious masses of mountains, so little to attract his attention and research, that the nature of the larger tracts of such country is but little understood. Mineral beds and veins are, for the most part, situated in hills, or eminences of more considerable elevation; to these, therefore, the attention of the miner and the mineralogist, as well as of the geologist, has hitherto been principally directed: the component masses of mountains are known the best. During the short period that mineralogy and geology have ranked among the sciences, they have made rapid advances: within the present century, considerable attention has been given to the exploring of some tracts of level country, which have amply paid the research. From the actual nature of these, we may reason by analogy of the rest.

In Europe, the principal tracts of low country are the eastern parts of England, the Netherlands, and the northern parts of France and Germany, and the whole of Poland. In Asia, the north-east parts of Russia, called the Steppes. In America, there are vast tracts of low land, through which the Mississippi and Missouri rivers take their course. The extent of the low lands of Africa are not ascertained.

From what is actually known, however, it may

be asserted that *the lowest and most level* parts of the earth, when penetrated to a very great depth, exhibit horizontal strata, composed of various substances, and containing, almost all of them, innumerable marine productions. Similar strata, with the same productions, compose the *hills* even to a considerable height. The shells are sometimes so numerous as to constitute the entire body of the stratum; they are often in so perfect a state of preservation as that their sharpest ridges are retained; they are found in elevations far above the level of the ocean, and in places to which the sea could not be conveyed by any existing cause: they are sometimes enclosed in loose sand, sometimes filled or penetrated by the hardest stones. Every part of the earth, every continent, and almost every island exhibits the same phenomenon.

It was once, long ago it is true, asserted that these remains of shells and other organized bodies, were merely the *sports of nature*; but it has been frequently found that the nicest and most scrupulous comparison of their forms, cannot detect the slightest difference between some of these shells, and the shells which still inhabit the sea. They have therefore once lived in the sea, and been deposited by it; the sea must consequently have rested in the places where the deposition has taken place. Hence it is evident that the basin or reservoir containing the sea has undergone

some change at least, either in extent, in situation, or in both.

The traces of revolutions become still more apparent and decisive if we ascend a little higher, and approach nearer to the foot of *great chains of mountains*; still many beds of shells are found, some even larger and more solid, quite as numerous and well preserved, but not of the same species as those found in less elevated regions. Here the strata that contain these shells, are of various degrees of inclination, and sometimes instead of being horizontal, as in plains and low hills, are even vertical. The strata of great chains of mountains, of whatever composed, or however placed, are laid open to view by means of vallies which time and violence have produced.

The diversity existing in the inclination of strata, clearly points out, in the estimation of some geologists, that by some means these have been broken and overturned.

The operation of an agent equal to the breaking up and overturning of the strata of mountains, and, if I may so say, to the destruction of rocks, and to the forming anew whole tracts of country which enclose the remains of organized bodies, was, it cannot be doubted, equal to the disruption of vast portions of continents, thereby forming islands: and it must, in all probability, have almost universally changed in appearance, and even in form, the surface of the globe.

It is beyond a doubt, that there have been many

catastrophes of the *same nature*, though not perhaps to the same extent. What has been the agent employed in the production of these catastrophes, is most obvious. It is not to be doubted that there have been successive irruptions and retreats of the sea; and it seems equally certain that the final result has been the universal depression of its level.

As we ascend to still higher points of elevation, and advance towards the lofty *summits of mountains*, the remains of marine animals, and that multitude of shells already spoken of, begin to grow rare, and, at length, disappear altogether. We arrive at strata of a very different nature, which contain *no vestige* of living creatures; nevertheless, certain circumstances observable in all these strata, in which not a trace of organic remains is to be found, have induced some geologists to suppose that their bare and rugged summits, though elevated far above the strata containing shells, have also been moved or overturned.

But though, by some, these rocks are not considered to be precisely in the place and position in which they were originally deposited, they are, nevertheless, considered by geologists to be older than all other rocks; because, as they contain no animal remains, it seems reasonable to suppose that they remain unaltered, in the same state as that in which they were created; and they have consequently received from geologists the name of *primitive rocks*.

Rocks of this description rise through others at various elevations in every quarter of the globe ; but in their greatest elevation, *primitive mountains* traverse our continent in various directions, rising above the clouds ; separating the basins of rivers from each other, and serving, by means of their perpetual snows, as reservoirs for feeding the springs ; and forming, in some measure, the skeleton, or, as it were, the rough frame-work of the earth.

I shall here recapitulate what has just now been said ; and shall afterwards proceed to its elucidation, by adducing such proofs, drawn from the observations of men who have made the phenomena of the globe their study, as may be consistent with the nature of the subject and our present object.

1. That the *lowest and most level* parts of the earth consist of horizontal strata, composed of various substances, many of them containing marine productions.

2. That *similar strata* are found *in hills* to a great height.

3. That shells are sometimes so numerous as to constitute entire strata.

4. That shells are found in elevations *far above the level of the sea*, and at heights to which the sea could not be raised by any existing cause.

5. That these shells once lived in the sea, and were deposited by it.

6. That shells continue to be found as we rise to the foot of *great chains of mountains*.

7. That, at this elevation, the strata, instead of being horizontal, as in plains, are of various degrees of inclination, and sometimes vertical.

8. That, from these and other circumstances, it is inferred that there have been frequent irruptions and retreats of the sea.

9. That, as we approach the *summits of lofty mountains*, the remains of marine animals and shells become rare, and even wholly disappear.

10. That their strata are wholly different, and contain no vestige of a living creature.

11. That these strata are, by some, considered not to be precisely in the place in which they were formed.

12. That, nevertheless, as they contain no vestige of animal remains, they are considered to be the oldest rocks, and therefore are called *primitive*.

The consideration of these points will naturally involve inquiries into others which may be termed subsidiary to them. In order, however, to form a complete outline of Geology, it will be necessary to add to these a number of other inquiries, more completely within the province of the mineralogist. Of these I shall now present an outline; and after attempting an elucidation of the foregoing, I shall, in like manner, proceed with these, and present to your notice the experience and obser-

vations respecting them, of men who have studiously investigated the phenomena of the globe.

13. That rocks, which, because they include no vestige of animal remains, are termed *primitive*, are of various kinds.

14. That rocks enclosing animal remains, are never found underneath, or supporting, those rocks which are termed *primitive*.

15. That some primitive rocks alternate with each other, but that granite is found beneath all others, and frequently overtops all the rest.

16. That rocks which include organic remains must have been formed *after* the shells they contain; and *therefore*, not being considered *primitive*, are by some termed *secondary rocks*: whence the terms, used by geologists of *primary* and *secondary formations*.

17. That there are many varieties of *secondary* rocks, each of which has received a geological appellation.

18. That there exists another class of substances not appropriately termed *rocks*; but which, being considered to be the *debris* or ruin of rocks, by their long exposure to the action of air and water, or both, are therefore termed *alluvial deposits*.

I now proceed to the illustration of the first position, viz.

That the lowest and most level parts of the earth consist of horizontal strata, composed of

various substances, many of them containing marine productions.

The illustrations necessary to this position will include many of the newest, most striking, and most important geological facts. These facts will prove, to a limited extent, another of our assertions, namely, that the catastrophes to which the surface of the globe has been subject have been numerous: and it will be shown that some of these have not been owing to irruptions of the sea, but to the agency of fresh water; and it will clearly appear that these irruptions of *fresh and salt water* have been *alternate*.

In order to make the whole of these circumstances more intelligible, it will be requisite a little to anticipate, by adverting, here, to some geological facts, which, according to our previous arrangement, would belong to another place. Geologists, who have had ample opportunity of examining mineral deposits in large and mountainous tracts of country, have satisfactorily ascertained that certain deposits are always found *beneath, never above*, certain other deposits. Investigation has proved that rocks which contain *no animal remains* are always found *beneath, never resting upon* those rocks which *do contain animal remains*: and also that those deposits which are termed *alluvial*, as gravel, sand, clay, &c. are never found *beneath* other rocks, but always *resting upon* them.

Thus much it seems necessary to premise of geo-

logical fact, previously to entering on a detail of the extraordinary circumstances which I shall adduce to prove the truth of our first position.

An investigation of the country surrounding Paris to a considerable extent, which by comparison may be termed low and level, has lately been accomplished by the eminent naturalist Cuvier, associated with the acute geologist Brongniart, and they have published a masterly delineation of the geological structure of the country.

Now as chalk makes its appearance on the edge of this district, and almost surrounds Paris, though at a considerable distance from it; and as the surrounding chalk is found to dip beneath the soil, or alluvial matter within the district, and has in many places, and at various depths, been discovered far beneath its surface, by the sinking of wells and pits; it is justly concluded that the soil on which Paris stands, and the surrounding country to a great extent, was actually deposited in a large hollow consisting of chalk, which, therefore, has been termed by Cuvier and Brongniart the *chalk basin of Paris*.

Immediately covering the chalk is found a small stratum of plastic clay, used in the manufactory of different kinds of pottery. On the plastic clay rests a deposit by *salt water*, thence termed a *marine formation*: above this rests a deposit by *fresh water*, thence termed a *fresh water formation*: next above is found a second *marine formation*: above it, a second *fresh water formation*; and upon

this rests an *alluvial deposit*. I now proceed to a more particular description of the nature of these five deposits:

1st *deposit*.—On the plastic clay covering the bottom of the *chalk basin*, it has been said that the *lower marine formation* rests. It consists of coarse limestone abounding in marine petrifactions: associated with it, is a series of strata in regular order, as marl, sandstone, &c. all of which enclose marine shells, many of them still retaining their pearly lustre. Occasionally, the space usually occupied by the limestone and the series of other strata, is entirely filled with siliceous limestone without shells, resting on the plastic clay and supporting the deposit about to be described.

2d *deposit*.—Upon the above *marine formation*, rests a deposit by *fresh water*. It consists of gypsum covered by a bed of white friable marl, enclosing petrified wood of the palm kind, and the remains of fishes and shells; the gypsum contains remains of *extinct* quadrupeds, birds, amphibious animals, fishes and shells, all of which are of *land or fresh water species*. This deposit is therefore called the *lower fresh water formation*.

In the *Gypsum*, Cuvier discovered the bones of 5 varieties of an extinct animal, which he calls the *paleotherium*, (signifying ancient large animal,) varying in size from a sheep to a horse; and the bones of 5 varieties of another extinct animal, which he calls the *enoplotherium*, (signifying beast without weapons; it had no canine teeth,) varying in size from the horse to the ass. Both these species

he considers to have been natives of the country over which Paris is now built.

He also found the bones of an unknown species of the *dog*, and of the *fox*; also of an *ichneumon* double the size of the living species. Nearly an entire skeleton of a quadruped of the genus *didelphis* was also found, but not belonging to any of the existing species, which are natives of America.

The fossil bones of birds, are not so readily known as those of other animals; but Cuvier describes some, found along with the bones of the extinct animals, as belonging to the pelican, the starling, and the quail tribes.

Of amphibious animals, the bones of the tortoise and the crocodile are recognised.

Of fossil fish, there are 5 varieties; most of them are allied to the present species of fresh water fish.

The shells all belong to fresh water fish.

3d deposit.—Above the beds of gypsum and marl, just described as containing the remains of *fresh water animals*, lie two beds of oyster-shells, separated by a bed of sand and sandstone without shells, from a bed of sand and sandstone, containing *marine shells*. These seem to have formed but one deposit *by salt water*.

The two beds of oyster shells are separated by a thin bed of white marl. The shells of the lower bed are numerous, thin, small, and brown. The upper bed of oyster-shells is very thick, and the shells are arranged as they are found in the ocean: the greater number of them are whole, and have both valves.

On the bed of sand and sandstone above described, rests a bed of clayey sand and marl, in which lies the *buhrstone*, or *millstone*. As it contains neither vegetable nor animal re-

mains, it seems not sufficiently characterized to be referred decidedly either to the preceding, which is a *marine* formation, or to the succeeding deposit, which is a *fresh water* formation.

4th *deposit*.—Above the buhrstone or millstone, lies a deposit of limestone and of siliceous substances, as flint, pitchstone, and jasper. The siliceous matter is sometimes allied in character to the millstone. But the essential character of the whole of the deposit, is, that it contains *fresh water* and *land shells*, nearly all of which belong to the genera now living in morasses. This formation extends 30 leagues to the south of Paris.

5th *deposit*.—Above the four deposits just described, as being alternately from fresh and salt water, lies the *alluvial* deposit, which appears to be by fresh water, and is composed of variously coloured sand, marl, and clay. It contains rolled stones of various kinds, but is most remarkable for its enclosing the remains of large organic bodies: in it are found great trunks of trees, bones of elephants, of oxen, rein-deer, and other large land animals.

This account of the contents of the *Paris chalk basin* contains important geological information. The space to which this investigation was limited, is, indeed, small, when compared with the surface of a globe of 25,000 miles in circumference, one-third of which is land. And if the investiga-

tion of so small a comparative spot, should not be deemed conclusive in regard to the probable nature of low and level countries in general, we have only to refer to the natural history of other parts of France and of other countries; of Germany, and of many tracts in the north of Europe, to be convinced, that at least in degree, the same effects have been almost universal.

But it was no longer ago than in the last year, that a geological investigation of certain districts in our country,* gave an evidence, in most points, perfectly coinciding with the observations of Cuvier, in regard to the chalk basin of Paris. The minutiae of this investigation were detailed in a very interesting communication to the Geological Society, since published in the 2d volume of its Transactions. At present we cannot do more than notice its results.

It appears that in this country there are two chalk basins, in a greater or lesser degree resembling that of Paris. One of them is called the *Isle of Wight basin*; the other, the *London basin*.

The *Isle of Wight basin* comprehends the district between Newport in that Island on the south, Southampton on the north, Brighton on the east, and Dorchester on the west. The strata which cover the chalk in this district, are individually and collectively of various thicknesses; occasionally only two or three of them are found.

* By T. Webster, M. G. S.

and sometimes only one of them; but there is one place on the southern edge of this basin, which proves beyond a doubt that the same causes which operated in the Paris basin, extended their influence to the Isle of Wight, and probably at the same period of time.

The place to which I allude is Headen Hill, forming a part of Alum Bay, near the western angle of the Isle of Wight. Of this hill, which is about 300 feet high, a natural section has been laid open, since its deposition, doubtless by the sea which borders it. Any one may, therefore, easily satisfy himself that it contains the same description of strata as have been found in the Paris basin, and precisely in the same order; that is to say, *alternate salt and fresh water deposits*, enclosing shells perfectly similar to those found in the Paris basin. Of the truth of this we have evidence. It is ascertained by a comparison of the shells taken from the corresponding deposits in both basins; and the animal remains so compared, are now deposited in the collection of the Geological Society in London.

The London basin begins at Deal, in Kent, and extends (not in a right line) by Canterbury to Gravesend; comprehending the whole of Kent north of that line, except the Isle of Thanet, which is of chalk. The edge of the basin from Gravesend crosses the Thames at Grays, extends to Purfleet, whence it crosses the Thames again, and passes nearly in a straight line to Guildford, and

from Guildford to a little west of Hungerford, in Berkshire, whence it turns nearly north-east, to Maidenhead, Eaton, Watford, Hertford, Stansted, and Thaxted; but its precise extent to the north of that place, is not ascertained. In a word, however, the chalk basin in which London is situated, is comprehended in an acute triangle, one of its longest sides extending from Hungerford somewhat to the north of Harwich, the other from Hungerford to Deal; its shorter side taking in the whole coast from the north of Harwich to Deal, with the exception of the Isle of Thanet.

The perfect coincidence of the London with the Paris and Isle of Wight basins, has not yet been quite satisfactorily ascertained, in regard to the alternate depositions by salt and fresh water. The stiff blue clay, which prevails to so great a depth almost every where round and beneath London, is *unquestionably a marine deposit*, as all its numerous animal remains are those of *sea animals*. This clay lies immediately under the fine bed of gravel on which London is built. The wells in London pass through this clay from 200 to 300 feet; at Tottenham, about 130; at Lord Spencer's, at Wimbledon, 430 feet; at Harrow on the Hill, 70 feet; at Primrose Hill, near Hampstead, 500 feet without success; and, except in the latter instance, all arrived at the same bed of white sand, from which the water rose.

By a paper lately read before the Royal Society, we find, that, at Brentford, they lately passed 200

feet through the stiff blue London clay, without arriving either at water or chalk: above the clay lies a stratum of sand, gravel, and water; over that another, of 1 to 9 feet of loam; then 7 feet of sandy gravel; and then above, 9 feet of loam. These strata *lying over the clay* contain a vast collection of the bones of elephants, both African and Indian, of the hippopotamus, the horns and jaws of oxen, the horns of deer, and both land and fresh water shells. These circumstances clearly prove these deposits to have resulted from *fresh water*; whereas the *clay on which they rest* contains only the remains of *sea animals*. It is therefore probable, that a perfect accordance between the contents of the London and Paris basins will hereafter be satisfactorily ascertained.

In the strata above, as well as in those below the chalk, in the north-eastern parts of England, there is a remarkable agreement in point of position, in several places; in some, one or more of the strata may be wanting, *but the order in which they lie seems never to be inverted*.

If, however, we would take a wider range for proofs of that *more general* existence of these catastrophes that have befallen the surface of the globe, we shall easily find conviction. We are speaking now only of those to which the *lower and most level* parts of the earth have been subjected.

The bones of a species of *rhinoceros*, different from either of the three species of Africa, Asia, and the Isle of Sumatra, have been dug out of the

alluvial soil near Canterbury; and since, in many places of Germany, France, and Italy. In Siberia, not only single bones and skulls, but the whole animal, with flesh and skin has been discovered.

In the alluvial soil of France and Italy, have been found the bones of an *hippopotamus*, allied to the two only species now known, inhabitants of Africa and Súmatra; as well as the bones of another animal not allied to these, and entirely different from any of the existing species of quadrupeds.

The *tapir* is an animal peculiar to South America, yet two fossil species have been found in Europe; the one small, the other gigantic: both have occurred in different parts of France, Germany, and Italy.

Of the *elephant*, the only existing species are those of Africa and Asia. One fossil species has been discovered, differing from each, but most nearly allied to the Asiatic. It is the *mammoth* of the Russians. The bones of the mammoth have been found in the alluvial soil near London, Northampton, Gloucester, Harwich, Norwich, in Salisbury plain, and in other places in England; they also occur in the north of Ireland; and in Sweden, Iceland, Russia, Poland, Germany, France, Holland, and Hungary, the bones and teeth have been met with in abundance. Its teeth have also been found in North and South America, and abundantly in Asiatic Russia. Pallas says, that from the Don to the Tchutskoiness, there is scarcely a river that does not afford the remains

of the mammoth, and that they are frequently imbedded in *alluvial soil, containing marine productions*; the skeletons are seldom complete, still more seldom is the fleshy part of the animal preserved: but an interesting instance of this has been described.*

Five species of an animal more nearly allied to the elephant than to any other living species, have also been discovered: it has been called the *mastodon*. The five species are all herbivorous, the largest is about the size of the elephant; but no living species of the mastodon has been discovered in any part of the world; the fossil remains were found in Europe and America.

All these fossil species of quadrupeds have occurred in *alluvial soil that covers the bottoms of vallies, or is spread over the surface of plains*; none have been found in high vallies: the bones of some were covered by *marine shells* and remains, others by *fresh water shells*; and as no remains of these animals have been seen in any solid rock, or in any high mountain, it seems probable that these animals fell victims to some of the latest catastrophes that have befallen the globe; and though some of them differ from their co-species now existing in the torrid zone, there seems some

* The following account of the singular discovery of the carcase of a mammoth, is given by professor Cuvier, as taken from a report in the supplement to the Journal du Nord, No. 80, by M. Adams, Adjunct member of the Academy of St. Petersburg.

^ In the year 1799, a Tungusian fisherman observed a strange shape-

reason for supposing them to have been inhabitants of the regions in which their bones are

less mass projecting from an ice-bank, near the mouth of a river in the north of Siberia, the nature of which he did not understand, and which was so high in the bank as to be beyond his reach. He went next year, observed the same object, which was then rather more disengaged from among the ice, but was still unable to conceive what it was. Towards the end of the following summer, 1801, he could distinctly see that it was the frozen carcase of an enormous animal, the entire flank of which and one of its tusks had become disengaged from the ice. In consequence of the ice beginning to melt earlier and to a greater degree than usual in 1803, the fifth year of this discovery, the enormous carcase became entirely disengaged, and fell down from the ice-crag on a sand-bank, forming part of the coast of the Arctic Ocean. In the month of March, of that year, the Tungusian carried away the two tusks, which he sold for the value of fifty rubles; and at this time a drawing was made of the animal, of which I possess a copy.

Two years afterwards, or in 1806, Mr. Adams went to examine this animal, which still remained on the sand-bank where it had fallen from the ice, but its body was then greatly mutilated. The *Jukuts* of the neighbourhood had taken away considerable quantities of its flesh to feed their dogs; and the wild animals, particularly the white bears, had also feasted on the carcase; yet the skeleton remained quite entire, except that one of the forelegs was gone. The entire spine, the pelvis, one shoulder-blade, and three legs, were still held together by their ligaments and by some remains of the skin; and the other shoulder-blade was found at a short distance. The head remained, covered by the dried skin, and the pupil of the eyes was still distinguishable. The brain also remained within the skull, but a good deal shrunk and dried up; and one of the ears was in excellent preservation, still retaining a tuft of strong bristly hair. The upper-lip was a good deal eaten away, and the under-lip was entirely gone, so that the teeth were distinctly seen. The animal was a male, and had a long mane on its neck.

The skin was extremely thick and heavy, and as much of it remained as required the exertions of ten men to carry away, which they did with considerable difficulty. More than thirty pounds weight of the hair and bristles of this animal were gathered from the wet sand-bank, having been trampled into the mud by the white bears, while devouring the carcase. Some of the hair was presented to our Museum of Natural History, by M. Targe, censor in the Lyceum of Charlemagne. It con-

found. The mammoth found whole in Siberia was too warmly clad for the torrid zone.

Additional proofs of the *extensiveness of these catastrophes*, both by salt and fresh water, may be found in the famous rock of Gibraltar, and at various places on the coast of the Mediterranean.

The rock of Gibraltar is principally limestone, and is traversed by fissures or hollowed into caves, which contain a peculiar compound mass, consisting of angular fragments of limestone, of bones, usually of ruminating animals, generally broken, never in skeletons; and of land shells, cemented together by a calcareous basis: the bones were for a long time thought to be those of monkeys, but Cuvier has, with his peculiar sagacity, considered some of them to belong to a species of antelope, others to a kind of mouse.

At *Cette*, the limestone includes bones like those of a rabbit; others similar to those of the field mouse, and of a bird of the sparrow tribe; the vertebræ of a serpent, together with the bones of some ruminating animal, and three various kinds of land shells.

sists of three distinct kinds. One of these is stiff black bristles, a foot or more in length; another is thinner bristles, or coarse flexible hair, of a reddish-brown colour; and the third is a coarse reddish-brown wool, which grew among the roots of the hair. These afford an undeniable proof, that this animal had belonged to a race of elephants inhabiting a cold region, with which we are now unacquainted, and by no means fitted to dwell in the torrid zone. It is also evident that this enormous animal must have been frozen up by the ice at the moment of its death.

At *Nice* and the *Antibes*, the rock also contained the bones of the horse.

At *Corsica*, the rock contains the bones of small quadrupeds, chiefly foreign to the place; as those of one inhabiting the coldest and wildest parts of *Siberia*; and enormous quantities of bones, some of which resemble those of the field mouse, and others those of the water rat.

In *Dalmatia*, the bones contained in the rock are principally like those of *Gibraltar*.

At *Concud*, in *Arragon*, the rock contains the bones of the ox, ass, a small kind of sheep, and many land and fresh water shells.

We have now, as it seems to me, satisfactorily proved our first position to a considerable extent, viz. 'That the lowest and most level parts of the earth, consist of horizontal strata, composed of various substances, many of which contain marine productions.'

LECTURE IV.

Organic Remains visible in hills and on the sides of elevated mountains—Strata of the Brocken mountain—Summits of lofty mountains contain no organic remains—Heights of mountains—Division of rocks into primitive, transition and fletz (or secondary) and alluvial—Their definitions.

On the last evening, the real objects of Geological inquiry were pointed out. It was shown that these consist of the *natural phenomena and facts* every where discoverable; and that without an ample and nice investigation of these, it is impossible for us ever to attain a reasonable knowledge of the earth: such a knowledge, I would say, as may be derived from some acquaintance with the component masses of its crust, and of their relative positions. It was also shown how incapable and absurd are the speculations of mere closet-philosophers; who, relying on their inventive powers, and on the extreme difficulty of contradicting their silly theories, indulged themselves in speculations scarcely more ridiculous than it would be to assert that the globe is an egg or an oyster.

During the last evening also, were laid down a series of *geological positions*, as they may be termed, which have been found to result from the

truly philosophical labours of men who have investigated the crust of the globe, perhaps to as great a height and to as great a depth, above and beneath the surface of the sea, as man can easily attain. These geological positions I proposed to illustrate by quoting the experience of the very men from whose labours they have resulted. The first of them, viz. 'That the *lowest* and *most level* parts of the earth, when penetrated to a great depth, exhibit nothing but *horizontal* strata, composed of various substances, and containing almost all of them innumerable marine productions,' was then elucidated by the investigation of the chalk basin of Paris, and of the chalk basins of London, and of the Isle of Wight: and not only the truth of the foregoing position was made clearly to appear, but also the novel and interesting facts, that in these basins there have been successive and alternate deposits from *salt* and *fresh* water; which is proved by the nature of their strata, and the organic remains they respectively contain. And it was further shown that these catastrophes, so fatal to animal life, have not been partial; inasmuch as they are readily and largely seen in almost every part of the European continent, and particularly on the coasts of the Mediterranean sea.

The object of our present inquiry into the nature of the constituent masses of the surface of the globe, is so extensive as not to admit of those immediate convictions of the truth of what may

be asserted respecting them, as might be desirable. I claim however this advantage ; I demand no assent to theory, for I will not broach a theory. I offer alone the results of inquiries among the *facts and phenomena of nature*, by men whose love of nature and of truth, has rendered their researches invaluable to science : researches amid regions always open to the investigations of the doubting or disbelieving. Amongst these men, let us remember that we have an Humboldt, a Werner, a Saussure, and a Cuvier. What but the love of truth and of science could have induced Humboldt to traverse whole continents, or to ascend the Andes more than 18,000 feet above the level of the sea ; or Werner, the great German geologist, to bestow his life in examining the rude and mountainous regions which surrounded him, and in teaching the results of his inquiries ? What but the love of truth and of science could have led Saussure to investigate every corner of the Alps, during twenty years ; or have induced Cuvier to bestow twenty-five years of his life in the study of comparative anatomy and osteology, with a view principally, if not solely, to the illustration of the nature of our globe ?

If from all that Humboldt, and Werner, and Saussure, and Cuvier, and many other intelligent geologists, have observed, in regard to the nature and respective positions of the great masses forming the crust of the earth, we were to select such parts as would immediately come in evidence of

the truth of the geological positions already submitted to your notice, scarcely fifty evenings would afford time sufficient for their recital. It is my object to bring the required evidence into the narrowest compass; I shall, therefore, select only such as may suffice to attain our object; taking care, at the same time, that it shall be of the most obvious kind that the nature of our inquiry will permit.

The positions already recited, begin with the *lowest and most level* parts of the earth; these we have considered. We now ascend a little, that is to say, to the *hills*; and after a short notice of their nature, shall rise to the consideration of the masses constituting *lofty mountains*; taking occasion, here, to present an outline of the divisions which the experience of geologists has taught them to make in rocks, as the component masses of the crust of the globe; showing the reasons for their division into primitive, transition and flœtz, (or secondary) and alluvial; and that of each of these there are many varieties.

The 2d position is, That strata containing shells are found in hills to a great height.

3d. That the shells are sometimes so numerous as to constitute entire strata.

4th. That shells are found in elevations *far above the level of the sea*, and at heights to which the sea could not be raised by any existing cause.

5th. That these shells once lived in the sea, and were deposited by it.

6th. That shells continue to be found as we rise to the foot of great chains of mountains.

These positions I purpose to consider together. The evidence of facts requisite for their support, and as proofs of their truth, need not detain us long.

The cliffs of the Isle of Sheppey, bordering our own river, the Thames, do not rise to any considerable height above the level of the water. They have, however, long been celebrated for the numerous organic remains found in them, a list of which, was published several years ago. This list has since been enriched by a gentleman now resident at Faversham, by the addition of above 700 different species of fossil fruits, berries, and ligneous seed vessels. Among the animal remains found in these cliffs, are several varieties of the crab, the jaws of crocodiles, and lobsters nearly whole. It deserves notice that all these remains, both vegetable and animal, are entirely impregnated with sulphuret of iron, or pyrites.

At Reading, in Berkshire, or rather in the elevated lands in its neighbourhood, are found considerable deposits of oyster-shells; it is remarkable that many of them are entire, having both their valves united; but the animal matter, or oyster, is perfectly decayed. These shells have not undergone the process of petrification; they are white and extremely brittle, and readily separate into

laminæ. At Touraine, in France, 100 miles from the sea, and about 9 feet under the surface, there is a bed of shells 9 leagues long, and about 20 feet thick. According to Ulloa, there are similar deposits in Peru. Such are likewise well known to exist in almost every part of Europe. In the neighbourhood of Bath, at a rather higher elevation, large tracts of limestone are found, consisting almost wholly of shells; which are also discoverable in great abundance in the Gloucestershire hills, and in other parts of England. In the cliffs near Whitby, a crocodile has been found; in those near Lyme, in Dorsetshire, their remains occur in considerable abundance; and in the chalk cliffs of Dover, some varieties of fossil shells. Still higher, in various parts of Germany, the fossil remains of fish are found in hills and rocks of various kinds of slate.

Let us, however, continue to ascend. Dolomieu found immense quantities of sea shells on the sides of Mount Ætna, 2000 feet above the level of the sea; and at the height of 2400 feet above the same level he found, in the mountain itself, regular strata of gray clay enclosing sea shells.

Some of the lower hills of the Appennine chain, contain many species of shells; they contain also the fossil bones of elephants, of rhinoceroses, of the hippopotamus, of whales, and of dolphins.

It is asserted by Cuvier, in regard to the shells which are found imbedded in some rocks, that
 ‘ A nice and scrupulous comparison of their forms,

of their contexture, and even of their composition, cannot detect the slightest difference between some of these shells, and the shells which still inhabit the sea ;' an assertion which perhaps no one, who has at all examined it, will presume to deny. Surely then, we have a right to assume that they once lived in the sea, and that they were deposited by it ; and if deposited by it, that the sea must have been once sufficiently elevated ; since we know no other cause adequate to the deposition of rocks enclosing sea shells, and to their deposition in regular strata. And when we take into consideration that Mont Perdu, which is the highest of the Appennines, and reaches an elevation of 11,000 feet above the level of the sea, encloses so immense a quantity of sea shells, as that some of its strata seem almost wholly composed of them, we shall at once assent to the position that shells are found in places to which the sea could not be conveyed by any existing cause.

Mont Perdu is by no means the only elevated mountain enclosing sea shells ; it is, however, one of the highest ; and we may readily infer that, if a mountain of so great elevation encloses them, they will be found in the strata at the foot of great chains of mountains.

According to Saussure, the Buet, a mountain which rises 10,000 feet above the level of the sea, contains no petrifications ; but the Salenche, the Mole, and others not exceeding 7000 feet, are

found to enclose petrifications, although they form a part of the same chain.

The Altain chain of primitive mountains in Siberia enclose no animal remains; but they are flanked on each side by a chain of hills which enclose marine shells.

The 7th position is, That at the foot of lofty mountains, the strata, instead of being horizontal, as in plains and low hills, are of various degrees of inclination, and sometimes vertical.

8th. That, from these and other circumstances, it is inferred that there have been successive irruptions and retreats of the sea.

9th. That as we approach the *summits of lofty mountains*, the remains of marine animals and shells become rare, and even wholly disappear.

10th. That their strata are wholly different, and contain no vestige of a living creature.

11th. That their strata are, by some, considered not to be precisely in the place in which they were formed.

12th. That, nevertheless, as they contain no vestige of animal remains, they are considered to be the oldest rocks, and, therefore, are called *primitive*.

In proof that the *strata*, as we approach the foot of lofty mountains, are not horizontal, as in low

hills, I shall present to your notice the section of a mountain in the Hartz Forest, in Germany, made from the description given by Werner himself. This mountain is called the Brocken, (see plate 3.) and rises to a considerable elevation, though it is not one of the highest mountains in Europe : but the result of the examination of its surrounding strata by Werner, is better evidence of the facts it discloses than we could perhaps obtain from any other source. The centre is of granite, which is, as it were, mantled all around by several successive and perfectly distinct strata ; the oldest next to the primitive rock ; and invariably, each succeeding stratum, being newer than the preceding, dips lower and lower, as we depart from the primitive rock around which they are successively deposited. The nature of the several component masses of this mountain, will be further noticed when we arrive at the consideration of the mineralogical differences existing in mountain rocks.

The successive deposition of these strata (for that they were successive will become more apparent where their geological differences shall have been pointed out) may, at least in degree, be urged in proof that there have been repeated irruptions and retreats of the sea : they will moreover evince that *the sea has not always deposited stony substances of the same kind*, inasmuch as these deposits are distinct, and even essentially different, in their natures : and the strata sur-

rounding this mountain may be brought in evidence, perfectly in agreement with numerous other observations, that the sea *has observed a regular succession as to the nature of its deposits.*

And, from these circumstances, it is reasonable to infer, that the *sea has undergone great changes in the nature of its fluid:* whence we may presume that there may have been *a succession of changes in the nature of the animals which inhabit it,* corresponding with the changes in the chemical nature of the sea.

That such changes have taken place in the natures of the inhabitants of the sea, we have abundant proof.

Not only do *the species and even genera of the shells change with the strata,* but it is generally the case that *the shells of ancient strata have forms peculiar to themselves;* that these forms gradually disappear; that they are not found in the strata recently deposited, nor in the existing seas: but the more recent strata enclose some species which the most experienced eye cannot distinguish from those *which now inhabit the ocean.*

The section of the Brocken mountain, shows the reason for our assertion, that, as we approach the summits of lofty mountains, the remains of marine animals and shells become rare, and even wholly disappear. It has already been stated that granite, which forms the centre and summit of this mountain, is considered to be the oldest of rocks, because it is found underneath all others, and fre-

quently rises through and overtops all other the constituent masses of mountains, as well as because it never contains animal remains; granite, in like manner, constitutes the highest parts of very many mountains of different elevations throughout the globe. But it is often found that, although in some countries the sides of very lofty mountains have been covered by succeeding strata to a very great height, yet, in other countries, primitive granite, of very inferior elevation, is exposed almost to the level of the sea without having any part of it covered by secondary deposition; and the fact seems to be that these secondary depositions have greatly varied in extent and in elevation.

We now come to the consideration of the summits of lofty mountains which contain no vestige of a living creature, and whose stratification, if it may be so called, differs from that of mountains of less elevation.

The summits of lofty mountains generally consist of one or two, and sometimes of alternating, deposits of some of the older rocks; which, for the reasons already given, have been termed *primitive*. Some of these rocks mostly assume one appearance; others have mostly an appearance wholly different: I say mostly, because there are but few rocks that always assume the same appearance in regard to stratification.

For instance, granite, and some other of the

older rocks, sometimes occur in regular and nearly horizontal strata; sometimes have no appearance of regular deposition, either horizontal or inclined; but the summits of lofty mountains, constituted of such rocks, seem composed of large and irregularly sized blocks, piled on each other without any appearance of order: while on the contrary, *gneiss*, another primitive rock, is almost invariably in horizontal strata.

This diversity in appearance is very considerably augmented in mountains consisting of alternate masses of primitive compounds, which are by no means rare; and as these rocks suffer, in different degrees, by long exposure to the action of the elements, this circumstance considerably contributes to increase the disorder of their rugged summits, which are described as appearing at a distance like the ruins of towers and of fortifications.

Whether these constituent masses are still in their original position, is a problem of no inconsiderable interest; nor can we wonder that able geologists should differ on the subject. Cuvier, on the one hand, considers that the very appearances of their summits, are so many proofs of *the violent manner in which they have been elevated*. He is of opinion that all the older strata of which the crust of the earth is composed, were originally in a horizontal position; and that they have been raised into their present highly inclined position, by

subsidiencies that have taken place over the whole earth.

On the other hand, Jameson (who is a rigid follower of the opinions of Werner, whence we may infer, that it is also Werner's opinion) believes, that the present inclined position of these strata is, in general, their original position; an opinion which he considers to be countenanced by the known connexion of strata, the phenomena of veins, the crystalline nature of the older rocks, and also by what he terms the great regularity in the direction of strata throughout the globe.

Since, therefore, two authorities so eminent have not yet decided the point, and since their opinions are directly opposed to each other, we must be content to await and to expect an agreement, drawn from yet further inquiries amid the phenomena presented by the grand features of mountain rocks.

But the researches and experience of many skilful geologists all unite in this: *that the strata composing the summits of lofty mountains, contain no vestige of animal or organic remains; they are, therefore, considered to be in their primitive state.* An approach towards these summits discovers that the sides are covered, or, like the Brocken mountain, mantled around to a very great elevation, by deposits enclosing sea shells and other organic remains. These are common in the lower Pyrennees, whose elevation does not exceed six or seven thousand feet above the sea:

according to Ulloa, however, shells have been found at the height of 14,220 feet above the sea on a mountain in Peru.

It is extremely difficult, it is even in most cases impossible, to ascertain the internal structure of large and elevated mountains : but if, on ascending them, it be found that their summits are crowned by certain rocks which are *known never to include shells or other organic remains, and which have never yet been found resting upon those rocks which do contain them* ; we have a right to conclude, from analogy, that the same rock which forms the summit, composes the mountain itself, ascending from the base to the summit through the centre ; and that the masses of rock surrounding it, even to a great elevation, were deposited after the creation of the central mountain.

We may, I say, conclude this to be the case from analogy, because, in certain districts of the European continent, the operations of the miner have occasionally disclosed the fact. In various parts of Germany some mines are situated in lofty mountains. For instance, in the Krivan mountain, there is a gold mine, 6954 feet above the sea ; and in the mountains of the Tyrol, a silver mine, 7512 French feet above the same level.

As we are now upon the subject of mountains, it may not be amiss here to introduce a sketch, exhibiting the comparative heights of some in various quarters of the globe, whose names are

best known to us, as most frequently occurring in the usual course of our reading: it is not intended to convey a representation of their actual form (see Plate 1.) To us, who live in a low and almost level country, Skiddaw and Helvellyn are objects of wonder and admiration; but when these, or Ben Nevis, which is the highest mountain in Britain, are compared with the majestic elevations of the European or the American Continents, they sink in our estimation into mere hills; and the great pyramid of Egypt, that wonder of ages, which is 315 feet in height, seems as nothing in point of bulk. The heights of two or three remarkable cities have been added.

The highest mountain in Europe is Mont Blanc, in Switzerland, which is 15,662 feet above the sea; but there are several of nearly the same height.

The highest mountain in Asia is Petcha, or Hamar, in Chinese Tartary, which is estimated at 15,000 feet above the plains of China; unless it be admitted, that the highest summit of the mountains of Thibet exceed it, which, according to Colonel Crauford, is about 25,000 above the sea.

In Africa, the highest mountains are supposed to be those of Geesh, which are estimated at 15,050 feet above the sea.

Chimborazo, the highest summit of the Andes, and the most elevated of the American continent, is 20,282 feet above the sea; but there are four-

teen other mountains on that continent, between 10,000 and 20,000 feet in elevation; three of which are volcanoes.

Heights of Mountains, &c.

		feet above the sea.
Britain	Ingelborough, Yorkshire	3000 ^p
	Ben Lomond, Scotland	3048
	Saddieback, Westmoreland	3240
	Helvellyn, Cumberland	3324
	Snowdon, Wales	3456
	Skiddaw, Cumberland	3530
	Schihallien, Scotland	3564
	Ben Nevis, Scotland	4350
Italy	Vesuvius	3900
France	Puy de Dome	5000
	Puy de Lanff	6200
	Plomb de Cantal	6300
Jamaica	Blue Mountains	7431
Germany	Lomnitz Peak	8640
	Kesmark Peak	8508
	Krivan	8300
Pyrennees	Canigou	9000
	Mont Perdu	11,000
Canary Islands	Peak of Teneriffe	11,424
Sicily	Ætna	10,032
Alps	Lake Lauzon, Mont Olan	6796
	Mont Titlis	10,818
	Schrekhorne	13,000
	Mont Rosa	15,000
	Mont Blanc	15,662
America	City of Mexico	7424
	City of Quito	9000
	Silver Mine of Jauricocha	15,500
	Tunguragao	16,170
	Cotopaxi	18,600
	Chimborazo	20,282

We have now arrived at that branch of our subject which may be termed *Mineral Geology*; which has for its object the natures and differences existing in the component masses of the earth.

These masses are, by Werner, divided into *primitive, transition, flatz, and alluvial*.

PRIMITIVE ROCKS never contain animal or other organic remains, and are never found to alternate with, or to rest upon those rocks, which enclose animal remains.

Primitive rocks are so named, because, in so far as we know, they are the oldest, and were the first formed. They have a crystalline appearance, and are, therefore, *chemical deposits*, principally composed of the *siliceous, argillaceous,** and *magnesian earths*. Granite, gneiss, mica-slate, clay-slate, primitive limestone, serpentine, porphyry, and sienite, are of this kind. Of these, granite is considered to be the oldest, and sienite the newest.

To primitive rocks succeeds another class, which Werner denominates **TRANSITION ROCKS**; these enclose organic remains of animals now inhabiting the seas, and are principally composed of *chemical deposits*; but amongst them *mechanical deposits* first make their appearance.†

* The earth called *alumine* forms the basis of common alum, whence it obtained that name; it enters largely into the composition of clays, whence it is termed the *argillaceous earth*, from the Latin *argilla*, clay.

† The difference between a chemical and a mechanical deposit may be thus explained.

Limestone, though it sometimes occurs among primitive rocks, first appears in considerable quantity among transition rocks; amongst which greywacke, greywacke-slate and transition limestone are the predominating rocks.

Still newer than transition rocks, is the extensive class of FLETZ ROCK; the *older* of these contain the remains of sea fish *approaching in character to the kinds now found in the ocean* and the *newer* of them contain *shells precisely the same as now exist in the sea*. Fløetz rocks are, for the most part, *mechanical deposits*. The principal among these are limestone and sandstone; to which may be added gypsum, salt, and great accumulation of inflammable matter in the state of coal.

A *chemical deposit* is the result of that law of nature which has been already explained, called *affinity*.

Pounded flint, which is nearly pure silex, with a certain proportion of potash and some other substances, when melted, forms a glass which is soluble in water. When thus dissolved, the mixture is called the *liquor of flints*. It is said that a quantity of this liquor was left in a bottle during eight years, by Professor Siegling; who found that the force of affinity acting upon the particles of silex in the liquor, caused them to be deposited in transparent crystals of quartz, hard enough to give fire with the steel. These crystals were therefore a real *chemical deposit*.

A *mechanical deposit* is effected without the agency of affinity. If, for instance, sand or clay be mixed up in water, it will be deposited without regularity, or intimate combination, merely by its own weight. If this deposit be left to dry, and to become hard, it will not break in any particular direction, nor will the particles of which it is composed adhere together very strongly. There is no regularity, it is a mere accidental or *mechanical deposit*.

L

Still newer is the class of ALLUVIAL ROCKS ; these contain the *shells of fish now existing in the seas, and the bones of large land animals* ; and are almost entirely composed of *mechanical deposits*. Sand, clay, loam, and brown coal, are the principal earthy masses that belong to this class.

Such is the arrangement which observation has taught the experienced Werner to make in rocks. Some other geologists, however, are of opinion, that the division made by him of rocks enclosing organic remains into two classes, transition and floetz, is unnecessary : they therefore term *all those rocks which contain organic remains*, excepting those called alluvial, SECONDARY ROCKS.

We learn from what has preceded

- 1st. That the older rocks are principally composed of the siliceous, argillaceous,* and magnesian earths.†
- 2nd. That the primitive parts of the crust of the earth are entirely *chemical* productions ; whereas, in the newer, we find a beginning, and in the still newer, an increasing quantity of *mechanical* depositions.
- 3d. That limestone occurs but sparingly in the primitive, more abundantly in the transition (or older secondary,) and in the floetz class (or newer secondary) in immense quantity.

* See last note but one.

† See notes to the descriptions of the nine oldest and most abundant of the primitive rocks.

4th. That in the earlier deposits we meet with no bituminous or saline matters, as coal or slate, but that these occur in great quantity in the newer formations.

But that part of our subject to which we are now arrived, viz. the consideration of the *nature of the individual rocks or compound masses*, which have been brought to light by the operations of the miner, or the researches of the geologist, opens to us a field of inquiry of such amazing extent, as to induce me to pause, and, for a moment, to consider the precise nature of the object we have in view.

This, if I rightly understand it, seems to be the acquisition of such *knowledge of the great outline of geological facts and phenomena*, as the researches of men eminent in science have enabled us to attain; avoiding, on the one hand, an attempt to enter with mineralogical exactness into the study of every geological compound, and on the other, the bare recital of a catalogue of geological names; neither of which would afford either interest or instruction.

In the consideration of the natures of individual rocks, are involved the remainder of our geological positions. We proceed to the

13th. That rocks which, because they conclude no vestige of animal remains, are termed *primitive*, are of various kinds.

14th. That rocks enclosing animal remains are never found underneath, or supporting, those rocks which are termed primitive.

15th. That some primitive rocks alternate with each other; but that granite is found beneath all others, and frequently overtops all the rest.

16th. That rocks which include organic remains must have been formed *after* the shells they contain, and, *therefore*, not being considered *primitive*, are, by some, termed *secondary rocks*: whence the terms used by geologists of *primary* and *secondary formations*.

17th. That there are many varieties of *secondary rocks*, each of which has received a geological appellation.

18th. That there exists another class of substances, not appropriately termed *rocks*, but which, being considered to be the *debris*, or ruin of rocks, by their long exposure to the action of air or water, or both, are therefore termed *alluvial deposits*.

I shall now invite your attention to that classification of rocks which the experience and observations of Werner have induced him to make.

PRIMITIVE ROCKS.

Granite	Porphyry
Gneiss	Sienite
Micaceous schistus	Topaz rock
Argillaceous schistus	Quartz rock
Primitive limestone	Primitive flinty slate
Primitive trap	Primitive gypsum
Serpentine	White stone

SECONDARY ROCKS;

or

TRANSITION ROCKS.	FLÆTZ ROCKS.
Transition limestone	Old red sandstone
Transition trap	Flætz limestone
Greywacke	Flætz gypsum
Transition flinty slate	Variegated sandstone
	Second flætz gypsum
	Shell limestone
	Third sandstone
	Rock salt
	Chalk
	Flætz trap
	Coal
	Newest flætz trap


ALLUVIAL DEPOSITES.

Sand, gravel, loam, clay, wood-coal, &c.

But before we proceed to examine these rocks individually, let us take some further notice of the interesting section of the Brocken Mountain, which throws much light on the subject of the relative situation of mountain masses in general. An outline of the nature of these masses will of course be comprehended in the succeeding description of individual rocks. The centre of the mountain is *granite*, (1,) on each side reposes another primitive rock, called *clayslate*, (2,) which, as well as all the succeeding strata, is found entirely to surround the *granite*. The two strata

next in succession (3, 4,) are by Werner termed transition rocks; the first being *limestone*, the next *greywacke* and *greywacke-slate*. The stratum resting on the latter is called by Werner, the *old red sandstone*, (5,) and is the oldest of what he terms the flötz rocks; the succeeding strata to (10,) inclusive, are also flötz rocks. On the old red sandstone reposes the *1st flötz limestone* (6); on it the *1st flötz gypsum* (7); then succeeds the *2d* or *variegated sandstone* (8); then the *2d* or *newer gypsum* (9); and lastly, the *2d limestone* (10). It is essential to be noticed, that I do not pretend to give the precise extent, dimension, or shape, of the granite forming the centre of this mountain, or of the several successive and incumbent strata; their general shape and position is all that is intended to be conveyed. Nor must we forget to observe that, in reality, all the strata incumbent on the granite, are less and less in elevation as they are more and more distant from it, each newer deposit being lower than that preceding it in point of age and situation. Nor must we fail to remark that these several strata, which may be supposed to stretch along through a considerable tract of country, would soon assume that position which determined Werner to give to the newer amongst them the term of flötz rocks; that is to say, they would be *flat*, which is the meaning of the word flötz. On the *2d limestone* (10) reposes the *alluvial deposit* (11); with the precise nature and extent of this I am not acquainted.

This section has so greatly the appearance of



Section of the Brocken Mountain Harz Forest Germany

Faint, illegible text, possibly bleed-through from the reverse side of the page. The text is arranged in approximately 20 horizontal lines. Some words are highlighted in red ink, including "Meyn" and "Lust".



order and of art, that it might be supposed to be the mere contrivance of some theorist to answer the particular objects of his invention: but we have no reason to suspect it. That Werner is in some sort a theorist, perhaps, can scarcely be denied; but it must, at the same time, be allowed, that his great object has been to develope a grand outline of the facts presented by nature, and that his theory is wholly built upon investigations, to which the great mining and mountainous district in which he resides is particularly favourable. A Cornish miner of observation and talent, but whose education and knowledge is principally confined to his art; who had never read, or perhaps even heard of a theory of the earth, lately told me that on examining a certain district in Wales, he was surprised to find on the side of a mountain, strata of various kinds in regular succession, and much more so to observe the *same strata in the same order* on the side of another mountain distant nearly 20 miles from the former. It was the more remarkable to him, because Cornwall is not a stratified country.

I have selected for your inspection the section of the Brocken mountain, as being a well authenticated example of remarkable order in its several deposits, and because the relative position of rocks, and the order in which they succeed or cover each other, forms a curious and interesting part of geological inquiry. If, however, we were to imagine that this forms a representation of the

deposits of mountain masses in the aggregate, we should err. Though granite frequently overtops other rocks, it is, perhaps, more frequently found, that other primitive rocks rest immediately *upon and above* it. Granite is sometimes observed to alternate with gneiss, and gneiss with micaceous schistus and clay-slate.

Mont Blanc, which is 15,680 feet high, and is the highest mountain in Europe, is of granite nearly to the summit; which is, according to Saussure, of argillaceous schistus. It is said that in the Andes, in South America, granite has not been seen higher than 11,500 feet above the sea. A mountain called Marno, in Portugal, is granite covered by clay-slate enclosing crystals of a mineral called the chiastolite. The same rock enclosing the same substance, forms the summit of Skiddaw in Cumberland, probably resting also on granite.

It is sometimes found that several of the primitive rocks rest upon the granite, and above these some of the transition or flœtz rocks; or, as they have been conjointly termed by some geologists, secondary rocks. Even alluvial matter is said to have been observed, covering the summits of elevated mountains.

But it has been objected to the rules laid down by Werner, in regard to the relative ages and positions of rocks, as forming mountain masses, that, as his observations have been chiefly confined to the country in which he resides, it is not rea-

sonable to adopt the result of researches with regard to one district, as obtaining in every other country. This objection may, in some degree, be well founded. The observations of other geologists in other countries may not be all in perfect accordance with the rules laid down by Werner: but this dissonance may be attributed in some measure to the yet imperfect state of the science, and to the want of precise definitions in regard to the characters of many rocks. Some, which have obtained the same appellations, are so different in appearance, that the most experienced eye alone can determine their general agreement in regard to character and composition; while, on the other hand, certain others have obtained different names in different countries. In this imperfect state of geological language, the knowledge and skill of the observer must always be had in consideration.

If Werner be actually a theorist, he is one of a superior order. He has extended his researches throughout the large and important district surrounding him. The relative age, deduced from the relative position, internal structure and contents of the great masses forming that mountainous district, seems to have been ascertained by him with a degree of certainty that defies the application of the term *theory* to his results. If he merit the name of theorist at all, it seems only to be in consequence of his assertion, or supposed assertion, (for hitherto his principal discoveries

have been communicated only by some of his pupils) that the same results will be found to prevail universally. It is certain that researches in almost every quarter of the globe, have tended in an astonishing degree to verify his opinions, that *order in regard to deposition is universally prevalent, and that this order is never inverted.*

It cannot, however, be denied, that theories built even upon researches into the phenomena presented by nature, have both advantages and disadvantages. They serve to induce research, inquiry, and discussion; which, when carried on with that temperate zeal which may be expected from men whose object is truth, will be sure to promote it. It were to be wished that the love of truth were thus prevalent; much needless and intemperate warmth would be averted; and were observers bent on this alone, as their ultimate object, science would be benefited by nicer and more candid geological description than can be expected from those, whose great object is the support of a favourite theory.

Inquiries into the nature of the globe contribute greatly to the advance of real science; they serve at once to amuse and to instruct, by affording ample materials for reflection; they force upon the mind an immovable conviction that the globe itself was called into existence by a Power, whose design and contrivance are every where manifest: a power whose immensity is unsearchable.

The few rocks that have been mentioned as composing the principal masses of primitive, transition, and floetz, or secundar rocks, and of alluvial deposites, comprehend, as it were, within their catalogue, and connected with their history, a variety of other rocks so vast and so numerous, and passing the one into the other by such imperceptible and nice gradations, that it may truly be said their study is the business of a life.

I now proceed to offer some remarks on the nature of individual rocks; beginning with the oldest of the primitive rocks, and taking occasion to notice some of their masses, remarkable for extent and elevation, as well as the metalliferous ores which principally abound in them.

GRANITE is a compound, granular, aggregated rock, composed of felspar, quartz, and mica, mostly in distinct crystals; sometimes the one, sometimes the other of these ingredients predominates, but most generally the felspar.*

But granite varies in respect to its granular character, from the very large to the very small; the large granular is considered to belong to the

* Quartz, felspar, and mica, enter into the composition of granite and gneiss, the two oldest of the primitive rocks. Quartz is composed of silex, with 2 or 3 per cent. of moisture or water. Felspar, of about 63 parts of silex, 17 of alumine, 3 of lime, 13 of potash, and 1 of oxide of iron. Mica, of about 47 parts of silex, 20 of alumine, 13 of potash, 15 of oxide of iron, and 2 of oxide of manganese.

oldest formations, the small and fine grained granite to the newest.

Werner considers that there are three sorts of granite: the primitive; a newer formation which traverses the primitive in veins, and the newest granite formation; this latter always rests on some of the older primitive rocks. It sometimes contains portions of other rocks.

The colour of granite is grayish white or reddish.

Granite sometimes contains other substances, as schorl, garnets, and tin ore: many metalliferous substances are found in veins that traverse it.

In the great mining field of Cornwall, both copper and tin occur in veins in granite in prodigious quantities.

When granite is exposed, it forms very high and steep cliffs; often also it appears as lofty and precipitous summits, denominated peaks; as is remarkably the case in Savoy, in Switzerland.

It occurs, as forming mountain masses, both *unstratified and stratified*; when in the former state, it presents large irregularly shaped masses, sometimes distinct globular concretions, as in Bohemia, the Hartz, and other places. Stratified granite occurs in Bohemia, Saxony, Switzerland, the Pyrennees, the Altain mountains, and in other countries. According to Jameson, 'Strata can only be formed by parallel seams, which have the same direction and extent throughout the mass.'

Granite is one of the most frequent and widely extended rocks: it forms the summits of the highest mountains in Scotland, in the Hartz, the Alps, in Bavaria, Bohemia, the Tyrol, and most countries in Europe. It forms a very considerable portion of the Uralian and Altain chain of mountains in-Siberia in Asia. In Africa, it forms the principal constituent part of the mountains of Upper Egypt, the Atlas mountains, and of the country about the Cape of Good Hope. In North America, it is found in New-York, Pennsylvania, and Virginia. In South America, it forms whole groups of the elevated Cordilleras; and a large tract of country extending from Cape Horn, which is the southern extremity of America.

It is remarkable, that in the mountainous regions of Peru, especially in the environs of volcanoes, no granite is found except in low situations, at the bottoms of valleys.

GNEISS is the next oldest of the primitive rocks, and like granite, it consists of felspar, quartz, and mica; it is generally so small-grained, and the mica for the most part prevails so much, that gneiss is mostly of a slaty structure.

It sometimes encloses other rocks, as granular limestone and porphyry. It contains occasionally beds of garnet, accompanied by lead ore and iron pyrites; and sometimes, though rarely, beds of slaty glance coal.

Mountains composed of gneiss are not so steep

as those composed of granite ; their summits are usually rounded.

There are few metals that do not occur in gneiss ; in which are situated the greatest part of the Saxon, Bohemian, and Salzburgian mines.

Gneiss mostly reposes on granite, but is sometimes incorporated in it, and sometimes alternates with it.

Ben Lomond, in Scotland, and Mount Rosa, in Italy, are almost wholly composed of gneiss, as well as the middle part of the Pyrennees. It abounds in Bohemia and Silesia, in Carinthia, in the Southern Alps, the Vosges, and in Scandinavia. It occurs also in Greece ; the mine works of the ancients, in the vicinity of Athens, are situated in it ; it is found also in Russia, in several districts of South America, in the Shetland Islands, and in many parts of the main land of Scotland.

MICACEOUS SCHISTUS is considered to be the next oldest of the primitive rocks. This is also called schistose mica, mica-slate, and glimmerschieffer. This rock is composed of mica and quartz,* and like gneiss, has a slaty structure. Garnets are so commonly found in it, as to be almost considered one of its constituent parts : occasionally it contains several other substances.

Micaceous schistus usually rests on gneiss, but is not commonly found on the summits of mountains ; nor does it often alternate with granite or

* For the analysis of mica and quartz, see note on granite.

gneiss; but generally reposes on the latter, round the sides of the primitive mountains, forming gentle acclivities. The summit of hills composed of micaceous schistus are round.

It contains some ores, as gold, iron, copper pyrites, and cobalt, with garnets and asbestos; but unlike gneiss, in which they occur in veins, it contains them in beds.

The most important mines of Sweden, as those of Dalecarlia and Fahlun; those of Roraas, in Norway; many in Hungary, Salzburg, Saxony, and Bohemia, are situated in micaceous schistus.

It is found in many parts of Scotland; the mountain Schihallien and the neighbouring country, are composed of it. Humboldt observed it in great quantities in South America.

ARGILLACEOUS SCHISTUS, or CLAY-SLATE, is a simple mountain rock,* and follows micaceous schistus in the great series of mountain rocks; but it sometimes contains thin layers of quartz, or more rarely of felspar. It also contains some other mineral substances, as schorl, garnet, and hornblende; and sometimes encloses other rocks.

Geologists enumerate four varieties of this rock,

* The three former rocks are called compound mountain rocks, being composed of two or three mineral substances, viz. quartz, felspar, and mica; but argillaceous schistus is called a simple mountain rock, because it is not so compounded. It consists of two or three elementary substances, as of siliceous, alumine, and oxide of iron. There are several varieties of this rock, which have not been accurately analysed.

one of which is of roofing slate; and eight varieties of rock peculiar to the clay-slate formation; amongst which is drawing slate. The impossibility of noticing all these varieties will be obvious, nor does it form a part of our present object.

Argillaceous schistus is one of the most metalliferous of the primitive rocks; it abundantly contains veins and beds of tin, lead, cobalt, silver, and copper; gold, and the ore of quicksilver are said also to occur in it. It is a very widely extended rock, and sometimes forms whole mountains and chains of mountains; but not of the most elevated: they have generally a gentle acclivity.

In the Highlands of Scotland, argillaceous schistus rests upon, and passes into, micaceous schistus. On the continent of Europe, it may be traced through a great extent of country; as in Saxony, Bohemia, Silesia, Franconia, Bavaria, the Alps of Switzerland, Austria, Hungary, and many other districts. It occurs in Pennsylvania in North America. It is said that nearly the whole country between Potosi and Lima, in South America, is composed of it.

Saussure found it on the summit of Mont Blanc. In Britain, it occurs largely in Cornwall, enclosing veins of tin and copper; and it forms the summit of Skiddaw, in Cumberland. The famous mountain of Potosi, in which are situated the great silver mines, consist entirely of clay-slate.

- PRIMITIVE LIMESTONE is the next in order of

primitive rocks. It is a simple mountain rock,* and is white, yellowish, grayish, greenish, or reddish white. Its structure is always granular. The oldest is the whitest and most granular.

It sometimes encloses quartz and mica; more rarely garnets, steatites, asbestos, and some other mineral substances.

Occasionally it occurs in distinct strata; or in beds, sometimes short and thick, sometimes so thick as to form whole mountains.

Primitive limestone is rarely found enclosed in the older rocks called gneiss and clay-slate, just described. As a rock, it contains various mineral ores in beds and veins; as those of lead, zinc, and iron, auriferous pyrites, and native gold.

Whole mountains in Stiria, Carinthia, Carniolia, and in the Pyrennees, are composed of it; as well as three in Switzerland 10,000 feet high. But primitive limestone in these situations, is generally in immense blocks, without any regularity in regard to size, dip, or direction; it is sometimes stratified, as at Altenburg, near the lake Neuenburg, and in some parts of Scotland.

* When lime is combined with the carbonic acid, the compound, when compact, is commonly called limestone; another variety is called calcareous spar, (calx being the Latin for lime,) and mountains of limestone are commonly called *calcareous* mountains.

Calcareous spar is composed of 57 parts of lime, and 43 of carbonic acid. The limestone of calcareous mountains yields about the same proportions.

The marble of Sutherland, in Scotland, is said to be particularly valuable. The promontory of Athos, in the Archipelago, is composed of primitive limestone; but the most extraordinary mass of it occurs in Spain: it is said that the mountain of Filabres consists of one block of white granular marble, 2000 feet high, and 3 miles in circumference, without any mixture of other earths or stones, and almost without a fissure.

PRIMITIVE TRAP: the rocks belonging to this formation are numerous; they are found in Scotland and Germany abundantly, and in some parts of Britain. *Trap* is a German word, signifying a *stair*. The rocks of this formation are termed trap rocks, because their exposure to the elements causes them to take the form of steps or stairs: they are very numerous; and, on that account, are less intelligible than most other primitive rocks.

Primitive trap is almost wholly composed of hornblende; which, in some varieties, is mingled with felspar, in others, with mica.*

When hornblende is mixed with felspar, it sometimes contains scales of mica, and is then called greenstone-slate; which occurs in great beds and mountain masses, and in Sweden forms

* Hornblende is a mineral which occurs massive occasionally, but mostly in crystals confusedly intersecting each other: its colour is dark green, approaching to black. It is composed of 42 parts of siliceous earth, 11 of lime, 12 of alumine, 32 of oxide of iron, and some water.

For the analysis of the occasional ingredients in primitive trap, felspar, and mica, see note on granite.

ranges of hills. Primitive trap is very metalliferous; celebrated mining districts in Sweden, Saxony, and Silesia, are situated in it.

SERPENTINE is considered the next in order of the primitive rocks: it is a simple mountain rock.*

Serpentine sometimes encloses other substances; and among them are asbestos, mica, and crystals of quartz and hornblende. But only one instance is known in which it is incidentally mingled in another rock: serpentine with limestone forms the precious stone denominated *verd antique*.

There appear to have been two or three formations of serpentine. Its first appearance is occasionally in gneiss, the oldest of rocks except granite, and afterwards in micaceous schistus. It never occurs in very distinct strata; but generally in shapeless masses and beds.

The ores of lead, silver, and copper, are found in serpentine, though not abundantly.

Serpentine rarely forms mountains, but is said to form the summit of Mont Rosa, which is principally of gneiss. In beds, it occurs in Silesia, Bohemia, Saxony, and in the Scottish islands; and it forms a large tract of country in Cornwall, of many miles in extent, in which occur veins of native copper, and some veins of the soap-stone, so abundantly used in potteries, which is sometimes accompanied by asbestos.

* Serpentine is composed of about 45 parts of siliceous earth, 30 of magnesia, 15 of alumina, with some oxide of iron, and water.

Some varieties of serpentine are exceedingly beautiful, and are turned to purposes of ornament. At Zobnitz, in Upper Saxony, several hundred persons are employed in quarrying, cutting, turning, and polishing the serpentine found in that neighbourhood; and the various articles into which it is made are carried all over Germany.

PORPHYRY is the next in age of primitive rocks, and is one of the most widely extended formations.

It is a compound rock, consisting of crystals of quartz or felspar, or both, imbedded in a basis which is considered to be of contemporaneous formation: this basis, in the older porphyries, is generally of hornstone or compact felspar: in the newer formations, the imbedding substance is generally of various kinds of clay.* Occasionally, porphyries contain portions of clay and agate, or chalcedony, and some other substances.

Porphyry sometimes contains the ores of gold, silver, lead, tin, iron, copper, and manganese; but these occur principally in the newer porphyries; always in veins, never in beds. The principal mines of Hungary are situated in porphyry.

Porphyry does not appear in distinct and well defined strata; but occurs in beds of great magni-

* For the analysis of the imbedded substances of porphyry, quartz, and felspar, see note on granite. It is difficult to give that of the imbedding substances which are numerous. It will suffice to say that they are principally composed of the siliceous and argillaceous earths.

tude, sometimes indeed forming mountain masses in various parts of the world.

It is found occasionally in Scotland and the Scottish isles. On the continent, it may be traced from Norway to the borders of the Black Sea. It appears in Sweden, Finland, the Hartz Forest, Saxony, Bohemia, Silesia, Salzburg, the Tyrol, Carinthia, Carniola, Greece, the islands of the Archipelago, Egypt, Siberia, and in North and South America.

SIENITE is the next oldest of the primitive rocks; it is less abundant than most, if not all, of the foregoing.

It is compound, and consists of hornblende and felspar; the felspar predominates: some varieties contain quartz and mica, and but little hornblende.* Sienite is very nearly allied to porphyry, and is equally metalliferous.

In the island of Cyprus it affords much copper; many of the important silver and gold mines of Hungary are situated in it.

Sienite occurs in Galloway, in Scotland: on the Continent, in the Electorate of Saxony, the forest of Thuringia, (where it abounds in iron,) Upper and Lower Hungary, Transylvania, the Isle of Cyprus, and Upper Egypt.

* For the analysis of hornblende, see note on primitive trap: and for the analysis of felspar, quartz, and mica, see that on granite.

According to Werner, there are yet five other primitive rocks, making in the whole fourteen: these are of far less abundant occurrence than the foregoing, and will not detain us long.

TOPAZ ROCK is composed of quartz, topaz, schorl, and a sort of clay: it is rare, having hitherto only been found near Auerbach, in Germany, where it forms a mountain mass of considerable extent.

QUARTZ ROCK is a simple mountain rock, occurring principally in veins and beds; but enclosing no metallic ores of any description. Sometimes it includes mica, and has then a slaty texture. Quartz occurs plentifully in certain mountains in Scotland, and in some of the Scottish isles. On the Continent, it appears in Saxony, Bohemia, Silesia, Bavaria, and other places. It is, however, said, that the mountain called Bultuc, one of the Altain chain, in Siberia, being 350 feet high, and 4800 broad and long, consists entirely of milk-white quartz; which sometimes forms spires on the tops of mountains, appearing like snow.

PRIMITIVE FLINTY SLATE occurs occasionally in beds, in some of the German and Saxon mountains, alternating in beds with clay-slate before described: it occurs also in veins. But this rock is scarcely known beyond those districts.

PRIMITIVE GYPSUM is perhaps the least important of all primitive rocks. It has hitherto been found only in the Alps in Switzerland, where it is granular; but, being mixed with mica and

clay-slate, the rock obtains a slaty structure : this structure is never to be observed in secondary gypsum, and therefore serves as a complete distinctive character.

WHITE STONE : the characteristic colour of this rock is white ; it is composed of a little mica and compact felspar, and has a slaty or granular structure : sometimes it contains garnets. Hitherto it has been principally found in Saxony and Moravia : a variety of it appears in the mountains of the South of Scotland.

We have now taken a cursory view of what, according to our present knowledge of the crust of the globe, are considered to be the *oldest rocks*. All these, though some of them be occasionally found mingled, or alternating in beds or strata with each other, are *crystalline deposits*, and are *absolutely without any trace of organic remains, either of plants or animals* : for this reason, these are supposed to be in the very state in which they were first deposited, and therefore have received the name of *primitive rocks*.

But the fourteen rocks of which we have taken a slight view, are not, by some geologists, supposed to form the whole catalogue of primitive rocks ; jasper, hornstone, pitchstone, hornblende-slate, and puddingstone, are, by some, added to it : some of these appear to be varieties of those we have described, and the characters of the rest, do not seem to be so essentially fixed, as forming moun-

tain masses, as to induce me to detain you by their description.

All rocks not included in the foregoing catalogue, (with the exception of those called alluvial,) have, by some geologists, been termed *secondary*, because they are found to contain more or less of organic remains: but it has been discovered that the four rocks found in immediate succession to the preceding fourteen, do not contain organic remains precisely of the same characters as the rest. For although the four rocks in question enclose some shells common to those immediately in succession to them, they also contain a variety of petrifications, distinct in their characters, called zoophites, or those animals which are considered as forming the first link in the chain of animated beings; *none of which are found in any of the succeeding rocks.*

In these four rocks, which contain the *first traces* of organic remains, also appear the first mechanical deposits; and Werner has termed them *transition rocks*, as connecting the *primitive* with the *newer rocks* containing abundance of the remains of plants and animals; these newer rocks he has called *flatz rocks*, (the word *flatz*, meaning *flat*,) because the position in which they are found is more flat than that of the primitive or transition rocks; and is frequently quite flat.

It has already been remarked that some geologists, who are of opinion that some of the distinctions made by Werner are not necessary, class

all those he has named transition and flœtz, under the general term of *secondary rocks*.

The four *transition rocks* are called transition limestone, transition trap, greywacke, and transition flinty slate.

TRANSITION LIMESTONE does not rise so high on the sides of mountains as primitive rocks; but it occurs in mountain masses, forming precipices, and narrow and deep vallies. It is not very metalliferous.

In beds, it occurs in Scotland, and in Derbyshire; on the Continent, in the Hartz Forest, and most other mining districts.

Of TRANSITION TRAP there are two varieties; one of them called amygdaloid, or, in Derbyshire, toadstone, forms immense beds with transition limestone in that county: it is found also in the mining districts in Germany.

GREYWACKE is a widely distributed and important rock; it presents the first appearance of a mechanical deposite.

It is described by Jameson as composed of grains of sand, connected together by a basis of clay slate.

Greywacke is uncommonly productive of metalliferous ores, both in beds and veins. Almost all the mines in the Hartz are situated in it, affording silver, copper, zinc, and lead. In Transylvania, greywacke is traversed by numerous small veins of gold.

This rock is also found in most European countries.

TRANSITION FLINTY SLATE is of small importance; it principally occurs in Bohemia, and in the lead hills in Scotland.

To the four transition rocks just described, succeed the flœtz rocks, which are twelve in number, according to Werner. It has just now been observed that the whole sixteen are frequently, if not commonly, denominated *secondary rocks*.

As the position of the flœtz rocks is mostly flat, we may correctly imagine them to exist as *twelve beds, deposited one above another*: and as they all contain animal or vegetable remains, I do not propose to give a minute description of each, but shall content myself with saying that they are found in most countries; in many, in prodigious quantities. Generally speaking, they contain few metalliferous substances; nevertheless some of them are of almost infinite value to man.

The first or oldest of these is called the OLD RED SANDSTONE, which, in point of age, succeeds the transition rocks just described.

On the red sandstone reposes the FIRST FLÆTZ LIMESTONE; which, in the Hartz Forest, yields copper and cobalt in veins.

On the first flœtz limestone lies the FIRST FLÆTZ GYPSUM; containing, in Italy, and on the banks of the Wolga, masses of sulphur of several hundred weight.

On the first gypsum, rests the **SECOND, or VARIEGATED SANDSTONE**; so called from its being marked with brown, red, and white stripes.

On the variegated sandstone lies the **SECOND or FLETZ GYPSUM**; and on this, the **SECOND or SHELL LIMESTONE**; remarkable for the immense quantity, as well as beauty and variety of organic remains, both of animals and plants, enclosed in it in various countries of Europe.

On the shelly limestone rests the **THIRD SANDSTONE**, remarkable in Bohemia for the picturesque scenery afforded by the numberless pillars and pyramids, single or joined together, two or three hundred feet high, over a large tract of country at Auerbach: in the neighbourhood also, caverns and grottos appear in the same sandstone, from which issue streams, that give rise to water-falls, and thus increase the beauty of the scene.

On this, which is the third floetz sandstone, rests the **ROCK SALT FORMATION**. Of this extensive and highly beneficial deposit, it is my intention to give a somewhat detailed account, particularly of our vast deposit in Cheshire.

To the rock salt formation, according to Werner, succeeds the **CHALK FORMATION**; the earthiness of which denotes the lateness of its origin.

To the chalk succeeds that which is called the **FLETZ TRAP FORMATION**; and over it the **INDEPENDENT COAL FORMATION**. As a description of some coal districts, particularly our own, would now occupy too much time, I purpose on a

future evening to give some account of this valuable deposit.

To the coal succeeds that which is by Werner termed the **NEWEST FLETZ TRAP FORMATION**; which is considered by many geologists to be entirely of volcanic origin.

These are the twelve formations, which, according to Werner, succeed each other in the order just described. By this we are not to suppose that we shall, in every place in which one of them is found, invariably find the rest: frequently many of them are wanting. In the sketch of the Brocken mountain, it will be noticed that the two primitive rocks, gneiss and micaceous schistus, are wanting. All, therefore, that we mean to say, in pointing out their successive formations, is, that if gneiss and micaceous schistus had occurred, they would have been situated between the granite and the limestone. So with the floetz rocks: if, for instance, we arrived at a bed of rock salt, we might possibly find it resting on a bed of the old red sandstone, although no fewer than six beds have just been described as *newer* than the old red sandstone, and *older* than the rock salt: but had these six beds occurred, they would have been found between them; and most probably in the order laid down.

But there is still another species of deposit of which we have to speak, improperly classed among rocks, which is invariably found above all other strata. I allude to **ALLUVIAL DEPOSITES**.

This class includes those substances that have been formed, and still are forming, in every quarter of the globe from previously existing rocks, owing to the action of air and water upon them. Alluvial deposits occur both in high and mountainous regions, and in flat country; filling up hollows in one, and forming plains in the other.

In *mountainous countries* they consist of rolled masses, principally of gravel or sand; sometimes irregularly heaped, sometimes forming beds, and containing fragments of ores, and some kinds of precious stones.

The disintegration of the surfaces of hills in some parts of Cornwall, affords an extensive illustration of the effects ascribed to the agency of air and water. Large deposits of tin ore, covered by the ruin of granite, are found in many places; and have afforded rich harvests to the miner. In one of the branches of Falmouth Harbour, the miner, after damming out the water, sunk through a bed of alluvial matter fifty feet in thickness; at the bottom of which was found a bed of rounded masses of tin ore, varying from two to ten feet in thickness; in which occasionally were mingled small grains of gold. It may be conceived that the ruin must have been great, when it is known that the profits reaped from the undertaking, amounted to at least £50,000. At the bottom was the solid rock, on which I have walked.

The diamond is found in alluvial soil in the

East Indies and South America, as well as the topaz and the hyacinth ; and gold on the coast of California.

In *low and flat countries*, sand, loam, clay, sulphur, bog iron ore, and beds of gravel, are found in alluvial soil.

In the loam and sand, great beds of bituminous wood sometimes occur ; of which the great underground forest in Prussia, yielding amber in abundance, is a remarkable instance.

The remains of plants and animals are found only in those rocks which are newer than those termed primitive, and rest upon them. Such remains however occur in very variable proportion, and even in some of the older secondary (or transition) rocks, no vestige of them is to be seen.

It has already been remarked, that in the transition rocks, which rest immediately on primitive rocks, occur the remains of animals called zoophites, which form the first link in the chain of animated beings. These seem to be most abundant in transition limestone ; in which, also, corals of different species are found, approaching very nearly, in external characters, to those now growing in tropical climates.

In transition trap and flinty slate, no organic remains are to be seen.

Greywacke and greywacke slate seldom contain petrifications ; in large tracts of country consisting of these rocks, not a trace is to be seen ; in others,

some few are found, both of animals and vegetables. The animal remains seem to be nearly the same as those found in the limestone. In greywacke it is said that the remains of animals of the serpent kind also occur: the vegetable petrifications appear like the stems and leaves of palm trees, and of reeds. Greywacke slate contains the remains of those remarkable corals, which are supposed to form the connexion between shells and corals.

The old red sandstone is the oldest of the floetz rocks: it contains but few petrifications, and these are principally of trunks and branches of trees, which seem to resemble those of tropical climates.

The first floetz limestone comprehends three or four varieties of rocks, in each of which animal and vegetable remains are found: some of these belong both to salt and fresh water; the remains of a large amphibious animal of the genus monitor have also occurred, which have been described by Cuvier.

All the other floetz rocks contain abundance of animal remains, except the gypsum and trap formation, in which they are rare: several of them also enclose vegetable remains. In the second floetz limestone, are found petrified fishes of various genera and species, and fossil amphibious animals; the stems and leaves of trees, and of flowers, as of the ranunculus. In the third floetz, or shell limestone, are found, together with prodigious quantities of shells, the fossil remains of

fishes and of birds. Chalk sometimes contains the teeth and bones of fishes; the remains of tortoises and crabs also occur in it.

In alluvial formations, petrifications are distributed, which often are so much rounded as to show that they have suffered by attrition: the organic remains are those both of fresh and of salt water, and also of land and of amphibious animals. The shells of oysters and of muscles are plentifully found; occasionally the teeth of sharks. In some places, the bones of the horse, the ox, and the stag occur, but differing from those of the living species; in others, the bones of elephants similar to those now inhabiting Asia and Africa, and of the rhinoceros and hippopotamus; in others again, the bones of several extinct species of the elephant; and of an elk, formerly an inhabitant of Ireland.

LECTURE V.

Of Mineral Veins—Of Salt Deposites—Of Coal Deposites—Of Volcanoes—Of the Deluge—Of the Internal structure of the Earth—Concluding Observations.

OF MINERAL VEINS.

IN treating of veins, we have a two-fold object. They merit our attention in respect to the extraordinary circumstances which attend them in all countries in which they occur; and also on account of their being the *chief mineral deposites*.

But mineral deposites are of two kinds; for metalliferous ores are largely found in beds, as well as in veins.

Mineral beds are for the most part horizontal; and are found both in primitive and secondary countries, of various elevation.

The ores of copper, iron, and lead occasionally occur together in beds in primitive mountains; and sometimes gold and silver are intermixed with them. Cobalt, and certain ores of mercury, also occur in beds. Almost all the metalliferous ores in the great mining district of Sweden, are in beds in primitive mountains. Lead, zinc, and iron ores occur abundantly in beds in secondary mountains.

In England, some ores are found in beds; but by far the greatest mineral deposites of this country are in veins: it is uniformly the case in Cornwall.

A vein may be described as a *fissure* that has been afterwards filled up with several different substances.

Humboldt observed a vein of calcareous spar 140 feet wide, traversing gneiss in the Alps of Switzerland. Jameson observed a vein of porphyry-slate traversing sandstone, in the Isle of Arran, nearly 160 feet wide; and in Scotland, veins of pitchstone and greenstone, from 10 to 100 feet wide. But these veins do not appear to have been what may be termed *metalliferous veins*; which, for the most part, are much narrower.

It is said, that in most *primitive* metalliferous mountains, veins extend but a few hundred fathoms in length, and that their width does not exceed two feet.

It has also been said, that a description of the veins of Cornwall would, generally speaking, suffice for those of almost every other country; and having heretofore given much attention to their actual state, I shall confine myself chiefly to them, and endeavour to give a general outline of their direction, length, depth, width, dip, and contents.

But in these respects, veins have not such an uniformity, as that the history of one would be an history of the rest: almost every vein has something peculiar in it; something to interest the geologist.

The *metalliferous veins* of Cornwall, that is, the veins producing copper and tin, which are the chief mineral productions of that county, run in

the direction of nearly *east and west*; they may vary a few points. There are, however, other veins, that *rarely contain any metallic substance*, which, for the most part, run *north and south*. These two facts are extremely curious.

Metalliferous veins may sometimes be traced along the surface of the earth, by a certain ochreous or rusty appearance; but this is not very common.

A vein may be said, in some sort, to resemble a deep cleft or crack in a field. This cleft, whatever might be its depth, must, of course, have a direction under ground: either it would be quite straight down, or it would have a slanting direction beneath the surface.

The veins of Cornwall scarcely ever take a direction quite straight down, or, in other words, quite at right angle with the horizon; but almost always either dip, or incline away from that angle.

So that the metalliferous veins which run east and west, dip or underlie either towards the north or south; and the non-metalliferous veins, which run north and south, dip either towards the east or west.

The length of no one vein in Cornwall has, as yet, been satisfactorily proved. Some of them have been traced two, or three, or even four miles; but no instance has occurred in which a vein has been known to stop. Nor has the miner ever yet seen the bottom of a vein. The length and depth to which veins extend, therefore, are not known.

There are several mines in Cornwall upwards of 1000 feet in depth from the surface, and two or three nearly, if not quite, 1300 feet deep.

Metalliferous veins differ exceedingly in regard to their width. A vein containing tin ore, in a mine called Whealan Coates, was only three inches wide, but was so rich as to be worth working; while another, in a mine called Relistian, was upwards of 30 feet wide, and was also very rich in tin. Some of the veins containing copper in Herland mine, did not exceed 6 inches in width; and so continued for a few fathoms, but eventually passed away east and west in mere strings, scarcely thicker than paper; but these veins yielded copper of a very rich quality. A copper vein in the next hill, varied from 12 to 24 feet in width, and was also very productive of copper.

But the generality of metalliferous veins, both of tin and copper, are from one to three feet in width: and these are preferred by the miner, because the ore they contain is generally less intermixed with other substances, than that of wider veins.

Hitherto we have been speaking principally of metalliferous veins. There is yet one circumstance, and a very important one, in regard to these, that we must not fail to notice. These veins are not *filled* with metalliferous ores. Were that the case, the miner would defeat his own object; because, as veins are very numerous, the quantity of tin and copper produced by them,

would be so great, as that the price he would obtain for the ore would not defray the charges attending his operations. The ores both of copper and tin principally occur in quantities which, though they may extend many fathoms every way, generally occupy, in point of fact, but a small comparative portion of the vein, and are therefore properly enough termed *bunches*.

A question here naturally arises. With what substances are the remaining parts of the veins filled up? These are occupied sometimes by rocks, or by stony or earthy substances of various descriptions; or by rubble, or refuse matter that seems to have resulted from the ruin of some parts of the neighbouring country. The non-metalliferous parts of a vein, of whatsoever composed, are commonly termed, by the miner, *deads*, because they yield him nothing. Sometimes, however, veins are found to have large empty spaces; but this, in Cornwall, is not common. Water is very abundant in veins, particularly in those rich in tin or copper. On a large mine it is not unusual to see two or three steam engines, for the purpose of drawing the water. These will raise and discharge into the neighbouring valley, at least 1000 gallons of water every minute, night and day.

The sides of metalliferous veins are generally very determinate; and are covered by a hard dark-coloured crust, called by the miner, the walls of the vein: and there generally runs down every

vein, a small vein of a whitish clayey substance, which sometimes adheres to one, sometimes to the other wall.

The ores of copper and the ore of tin (for of tin there is but one description of ore, while there are many of copper) do not often occur together in the same vein to any great depth beneath the surface.

At about 80 or 100 feet under the surface, the first traces of copper or tin are usually found; rarely nearer to it than 80 feet. But when at last they are found, it is not to be understood that these ores consist of one close and compact mass: on the contrary, they are generally mingled with other substances, such as lead ore, iron pyrites, the ore of zinc, accompanied by fluor spar and quartz, &c. These are in some cases loose in the vein; in others, they are hard, and attached to one or to both sides of it.

Sometimes the ores *both* of tin and copper are found thus circumstanced *together* in the same vein; and when so found, it generally happens that all trace of tin is soon lost.

If tin be first discovered, even without a trace of copper, it is not unusual, that in the course of sinking 80 or 100 feet more, all trace of it is lost, and copper only is found. The vein of course was at first called a tin vein; but afterwards became only a copper vein: and many of the most productive mines in Cornwall have been exactly so circumstanced. Nevertheless, in some veins,

tin continues to be found to the great depth of nearly 1000 feet beneath the surface, almost without a trace of copper.

But if, instead of tin, copper be first discovered at the depth of 80 or 100 feet, it seldom or ever happens that tin is found below it in the same vein.

In one or two of the deepest mines in the county, both copper and tin have continued down together, in the same vein, to the greatest depth at which it has been seen by the miner; sometimes one prevailing, sometimes the other.

It has been stated, that the *tin and copper veins run nearly east and west*; but that the *veins which run nearly north and south, scarcely ever contain a trace of tin or copper*: in some few instances, they have been found to contain the ores of silver, lead, cobalt, and iron; others have produced antimony.

These north and south veins are usually filled by quartz, or a whitish or bluish clayey substance, or an ochreous substance; and sometimes by all three. When a vein of this description meets with a vein containing tin or copper, it *passes through the tin or copper vein*; and sometimes, as it were, splits it into numerous little branches; the north and south vein continuing its course straight forward without interruption.

Not only is this curious effect produced, but also another of a much more extraordinary nature. In searching for the tin or copper vein on the

other side of the north and south vein, it sometimes cannot be found for a length of time, nor without much labour and expense: forty years have been spent in such a search.

For, instead of continuing its course, instances have been known in which the tin or copper vein has been again found 120, or even 450 feet, north or south of that part of it, on the other side of the north and south vein.

North and south veins vary in width from one inch, to ten or twelve feet; but, whatever be their width, they always divide tin or copper veins, and generally alter their course; or, in the language of the miner, *heave* them out of their course.


In some parts of the mining districts of Cornwall, metalliferous veins are so numerous, that with the miner, the question is not where a vein can be found, but where he will be most likely to meet with one productive of copper or tin. Years of labour, and large sums of money, are often expended in vain, because there is no circumstance by which he can determine, with certainty, that his efforts will be successful.

There are many mines through which several veins of copper and tin take their course, very near to each other.


If a copper vein meets with a tin vein, it is universally the case, that *the copper vein passes through that of tin*, and generally heaves it out of its course, greatly to the inconvenience and loss of the miner, who is often puzzled to find it again.



Veins in TIN CROFT Copper & Tin Mine



Veins in the PINK Mine



There are still other, and, if I may so say, subordinate veins found in Cornwall. The explanation of their nature and effects would trespass too greatly on your time: they rarely contain any metalliferous substance, but they occasion prodigious vexation and expense to the miner.

In elucidation of what has just been said of the phenomena attending the veins in Cornwall, I shall offer for your inspection a sketch of those which actually occur in two mines called *Tin Croft* and the *Pink*, (see plate 4.) Let us suppose that the upper square figure represents the hill in which Tin Croft mine is situated: it is not an exact representation of it, because that hill has a gentle declivity on every side; but for the sake of familiar illustration, I have supposed it to be cut away on the north, the south, the east, and the west sides, and that the soil is taken from the surface down to the rock, in order more clearly to show the run of the several veins upon it, and their directions beneath it.

All these veins are found in Tin Croft mine, in less than half a mile from north to south. The upper part of the square figure represents the field in which the mine is situated. We shall observe that there are three veins of copper, (coloured red;) three of tin, (black;) and one yielding both copper and tin, (red and black.) These veins run on the surface east and west, and are intersected by a vein running north and

south (bluish) which is not metalliferous; and we shall further observe, that the copper and tin veins were not merely intersected, but also, that the parts of them on the western side of the north and south vein, were 'heaved' out of their regular course towards the south.

Let us remark the downward direction of the copper and tin veins; not one of them runs straight down; they dip or underlie more or less, either towards the north or south. Two of the copper veins intersect one of the tin veins, and pass through it, without altering its direction.

It has been said, that the miner has never seen the bottom of a vein. Three of the veins in Tin Croft seem to stop; the fact is, that the miner did not pursue these veins to so great a depth as the others; their direction beneath where they seem to stop, is not exhibited, because it is not known.

From one of the copper veins which underlies a little towards the south, two branches go off, underlying much quicker towards the north: the veins of Cornwall rarely branch off in this manner.

The section of the Pink mine is a remarkable instance in proof of the assertion that *copper veins meeting with those of tin always divide, and pass through, and mostly 'heave' them.*

Let us first notice the summit of the lower square figure, (pl. 4.) which represents the run of the veins on the surface of the Pink mine; in which there were four east and west, or metalli-

ferous veins, two being of copper (coloured red,) and two of tin (black.) These veins were intersected by a north and south vein (bluish) that was not metalliferous; which heaved the parts of the copper and tin veins on the western side of it, more towards the north than are those parts of them on the eastern side of it.

The tin vein ran near the southern extremity of the mine, underlying in its downward direction greatly towards the north. It will be noticed that the copper veins underlie in the contrary direction, that is, towards the south. One of them, meeting with the tin vein in its course, interrupted it, and 'heaved' that part of the tin vein on the south of the point of intersection, twenty-four fathoms nearer the surface. It was afterwards found that the tin vein was again interrupted by another copper vein, and again heaved towards the surface, though only about ten fathoms: a third time it was cut through by still another copper vein, which does not appear on the surface of the mine, and again heaved, though less than before, towards the surface.

There are yet some important points respecting mineral veins, on which I purpose saying a few words. I mean in regard to the probable manner in which they were formed, and by what means they were filled.

They who contend that the great masses of the globe are altogether what they now are, *through*

the agency of fire, assert that veins were formed by the contractions which took place in those masses while cooling, and that they were filled from below: in other words, that the contents of veins were protruded into them from the internal parts of the globe, by the agency of fire.

Those who, on the contrary, contend that the great masses of the globe are what they now are *through the agency of water*, assert that mineral veins were originally open fissures or rents, caused by the subsiding of the great masses of the globe; and that these fissures were filled from above, receiving into them the metals which formed a part of a great chaotic fluid.

I am not now about to enter into an examination of the comparative merits of these two doctrines, but shall probably hereafter say a few words on this part of the subject.

I cannot, however, pass by one or two observations in regard to the relative ages of veins.

It has been said that copper veins, meeting in their course with those of tin, always divide and pass through them. This seems clearly to show that tin veins are the oldest, or they could not have been so divided.

It has also been said that the north and south, or non-metalliferous veins, always divide the veins of copper as well as those of tin. This clearly shows that the tin and copper veins were both older than the north and south veins, or they could not have been so divided by them.

But other veins, not containing any metallic substance, are occasionally found ; which, as they divide and pass through every one of the fore-mentioned, are therefore of still later formation.



OF SALT DEPOSITES.

It was stated during the last evening, that de-posites of salt are principally found amongst those which are by Werner termed the *flötz* or flat rocks ; but which, by other geologists, are ranked among those called secondary rocks.

Clay, sandstone, and gypsum, almost invariably accompany rock salt, either above or below it ; sometimes both above and below it.

The countries in which large depositories of salt are found, are for the most part flat ; they do not often exceed that elevation which is termed hilly.

In Germany, but few instances of the rock salt formation occur ; but it is said that an uncommonly great deposition of it may be traced with little interruption, from the Black Sea nearly to the Alps. It abounds in Spain ; but is not very common in Russia, or generally in northern countries. Nevertheless, there are said to be two whole mountains in Astracan entirely composed of it. It is abundant in Persia ; the isle of Ormus in the Persian Gulf almost wholly consists of rock salt. Whole mountains of it also occur in Tunis

and Algiers, in Africa. It is found in New South Wales; and not long since a mountain of salt, of an immense height, was discovered near the Missouri river in America, eighty miles long and forty-five miles wide, the surface of which is barely covered with earth; neither tree nor shrub is growing upon it.

But many countries are nearly without salt. At Delhi and Agra, the capitals of Hindustan, its price is *2s. 6d.* per pound: and it is said to be so scarce in the interior of that country, west of Thibet, that the natives use cakes of salt, sealed up and bearing the stamp of their prince, as money.

Perhaps the most extensive deposition of rock salt in the world, occurs in Wielitska, near Cracow in Poland, at the northern extremity of a branch of the Carpathian mountains. It has been worked as a mine since the year 1251, and its excavations are said to extend more than a league from east to west. The salt is of an iron gray colour, in which are found cubes of a pure white.

This mine was visited by our countryman Wraxall; from whose account of it some idea of its vastness may be gathered. He says, 'After being let down by a rope two hundred and thirty feet, our conductors led us through galleries, which for loftiness and breadth seemed rather to resemble the avenues to some subterraneous palace, than passages cut in a mine: they were perfectly dry in every part, and terminated in two chapels

composed entirely of salt, hewn out of the solid mass. The images which adorned the altars, as well as the pillars and ornaments, were all of the same transparent material; the points and spars of which, reflecting the rays of light from the lamps which the guides held in their hands, produced an effect equally novel and beautiful. Descending lower into the earth, by means of ladders, I found myself in an immense hall or cavern of salt, many hundred feet in length, breadth, and dimensions, the floor and sides of which were cut with exact regularity. One thousand persons might dine in it without inconvenience, and the eye in vain attempted to define or trace its limits. Nothing could be more sublime than this vast subterraneous apartment, illuminated by flambeaux, which faintly discovered its prodigious magnitude, and left the imagination at liberty to enlarge it indefinitely.'

Hitherto we have not mentioned the deposits of salt, and the salt or brine springs, which are so abundantly found in our own country. A description of some of these is my principal object. The chiefest are those of Droitwich, in Worcestershire, and of Northwich, in Cheshire, which are the most productive of all.

I proceed first to the *brine springs* at Droitwich.

These springs are said to be mentioned in the Domesday Survey, which was finished in 1087.

The prevailing rock around Droitwich is a

brownish red sandstone, considered to be the old red sandstone of Werner.

At this place the brine springs are four in number, all situated within a square furlong; and as no new pit has been sunk within the last thirty years, we have not a very accurate account of the strata through which they passed, before they arrived at the brine. From the account given by Dr. Nash, in his history of Worcestershire, we gather these facts: that four pits were variously sunk through from thirty to fifty-five feet of soil and rock, when they arrived at a stratum of gypsum, varying from 102 to 150 feet in thickness: on passing through this, they suddenly arrived at the salt brine, which, immediately on their arriving at it, rose quickly to the surface, and overflowed. In each of the pits the brine was twenty-two inches in depth: in each also it was ascertained that it was immediately resting on a body of rock salt. Into this rock salt they bored two feet and a half, without passing through it; the brine being their object.

From this account, the principal information we gather, is, that the water of these springs is impregnated with salt, by a body of rock salt; and that, as the brine rises perhaps 180 feet to the surface, and overflows, the source of these springs must be situated in much higher ground than that in which the pits are sunk. The brine is perfectly limpid, and contains about one-third its weight of salt.

The quantity of brine which issues from these four pits is immense. That which is used, bears but a small proportion to that which runs to waste: nevertheless, the quantity of salt annually made from these four pits is about 16,000 tons; two-thirds of this are consumed in England, and pay a duty of about £320,000 per annum. The market price of the salt is £31 per ton, £30 of which is duty.

We come now to the great beds of rock salt at Northwich in Cheshire.

These beds are known to extend one mile and a half, north-east and south-west, and are upwards of three quarters of a mile wide: there are two beds, lying one beneath the other.

The strata above the upper bed, consist of gypsum, and of alternating beds of variously coloured marl, red, blue, and brown; some of them are so porous, as that it has been ascertained that 360 gallons of water rise through them in a minute; a circumstance that greatly impedes the sinking of the pits. It is remarkable, but it is well ascertained, that the various strata above the upper bed of rock salt contain no marine fossils. These strata are from 105 to 120 feet thick; they repose on the first bed of salt, which is from sixty to ninety feet thick: between the first and second beds of salt lies a stratum of indurated marl, thirty-six to forty feet in thickness. So that the surface of the second or lower bed of rock salt is about

220 feet from the surface of the land. Into this second bed of salt they have sunk 132 feet, without having found the bottom of it.

The salt of these mines is, for the most part, of a reddish hue, arising from some admixture of iron; and it is generally so hard, that the blast by gunpowder is employed in breaking it down. The lower part of the lower bed is the purest: and in it there are considerable cavities, about 16 feet in height; in which, occasionally, pillars of salt are left, six or eight yards square, which form the supports of the roof. The cavities are worked into aisles or streets; which, when illuminated by candles fixed to the sides of the rock, give a brilliancy of effect that is singularly striking; and, it is said, almost appear to realize the magic palaces of the eastern poets.

Some idea of the vast magnitude of the Cheshire salt deposits may be formed, when it is mentioned that its many mines yield 16,000 tons for home consumption annually, and that 140,000 tons more are annually exported from Liverpool.

We come now to the consideration of the means by which these vast formations of rock salt were deposited. It must be obvious, that all that can be said will amount to no more than theory: a theory which presents some objections, while its basis seems reasonable in itself.

The Cheshire salt beds occupy vallies surrounded by hills of secondary formation; and the

upper surface of the upper bed of salt is about 40 feet below low-water mark at Liverpool. The numerous facts already adduced, have led us decidedly to adopt the belief that the sea must have stood at an elevation greatly above the general level of the earth. Is it not, therefore, reasonable to presume that these beds of salt were deposited by the sea; and that the beds of clay lying between and above them resulted from the ruin of rocks?

The arguments in support of this theory are,

That the upper surface of these deposits of salt, are 40 feet under low-water mark.

That, in the beds of marl, it is not unusual to find fragments of the older rocks; such as large portions of granite, showing marks of attrition.

That these deposits of rock salt contain some salts, as the sulphate of soda, which also is found in the waters of the ocean.

The principal argument against this theory is, that no sea shells or weeds are found above or below these beds.

But, perhaps, we ought to take into consideration that there are beds of salt near Salzburg in Austria, which are stated by Von Buch to be 2975 feet above the sea; and that the salt mines in the Tyrol are yet higher.

We have, however, heretofore produced evidence that the sea has been at much greater elevations than this; and has deposited almost entire strata of sea-shells, at such elevations.

OF COAL DEPOSITES.

Deposites of coal are not only of great importance, but are far more abundant and more general than they are commonly supposed to be.

Coal, in greater or less quantity, and of different qualities, is found in most countries: in Holland, Germany, Saxony, Portugal, Switzerland, Sweden, China, Japan, New Holland, and in North and South America. Buffon states, that in his time there were no fewer than 400 collieries worked in France. The deepest coal mine in the world is near Namur; it is stated to be 2400 feet, or nearly half a mile, in depth.

Coal is found at various elevations; but almost all the greatest deposits are in low situations; where it occurs in beds lying over each other, of various thicknesses, having between them one or more earthy deposits, or beds of stony matter. It is remarkable that though these beds of coal are nearly horizontal, they are not quite flat, but generally dip near the middle, where they are found thicker than at the sides; so that a section would give the idea of the form of a boat.

Both in this country and in others, coal is found at elevations much above the sea; and at one place in France, the strata, instead of being, as is usual, nearly horizontal, are nearly vertical. I have lately been informed that there is a considerable deposit of coal in the immediate

neighbourhood of the great silver mine of Jauricocha in Peru, which is about three miles above the level of the sea.

But it is considered that there are three formations of coal. I shall begin with the newest, on account of its being of the least importance.

The *newest coal formation* occurs in alluvial soils. In this, the strata of coal are not parallel with each other; and the earthy strata that are found with it are those of sand, clay, and gravel.

The newest coal consists almost exclusively of earthy brown coal, and bitumenized wood, of which there is a considerable deposit near Exeter.

The *coal formation, next* in point of age, occurs in that deposition to which Werner has given the name of the newest floetz trap; the result, as he conceives, of deluges. In this the coal is generally covered with clay, or basalt, in which are found neither vegetable impressions nor animal remains. The strata are not so numerous as in the formation presently to be described, nor are they so perfectly parallel with each other. The chief collieries of Scotland, of the central part of France, and of Bohemia, are of this description; these yield principally the varieties termed pitch coal and moor coal, not often slate coal.

The next, or *oldest*, is called the *independent coal formation*, because the individual depositions or beds, not being connected, are independent of each other. This formation consists of extensive and remarkably parallel strata of coal, covered

by strata of indurated clay called shale, containing the impressions of vegetables, and sometimes the remains of fresh water shell-fish. The shale is always wanting in the *newer* deposits of coal already described, but always accompanies the *oldest*. The great coal deposits of our own country, are principally of this latter description; and in some of these there are as many as twenty beds of coal, varying in thickness from 6 inches to 6 or 8 feet. Between the beds or strata of coal, is one or more beds or strata of various coloured sandstone, clay, bituminous shale, or rubble stone, (called by the miners rotten stone) or argillaceous iron ore, or of secondary limestone. This formation of coal is also plentifully found in some of the countries already enumerated; and two circumstances are worthy of notice. The first, that the strata occurring above and below this formation, are in all countries very much alike, if not absolutely the same: the second, that although the shale, already described as lying above the coal, contains impressions of vegetables, and the remains of fresh water fish; it is remarkable that in every country the strata of various substances which lie between the strata of coal, scarcely ever contain any vegetable impressions or organic remains.

During the last evening, in speaking of what may be termed *the order every where observable in the disposition of* great masses which form the crust of the globe, it was stated that deposits

of coal are principally found resting upon secondary rocks : and this is proved to be the fact in respect of our own great deposits, as well as those of other countries.

Our own deposits are our immediate objects.

Three extensive collieries in Flintshire in North Wales, those of Glamorganshire in South Wales, of Coalbrookdale in Shropshire, and of Kingwood near Bristol, all commence in the immediate vicinity of secondary limestone ; and the still more extensive deposits in the North of England, at Newcastle and Whitehaven, rest upon secondary freestone or sandstone, abundantly used as grindstones.

These two extensive collieries of Newcastle and Whitehaven, situated, the one on the north-east and the other on the north-west coast, it is now confidently believed, from the great similarity existing in their strata, form but one deposit, consisting of many strata, which extend directly across the island from one place to the other ; and even far beyond each, beneath the sea. At Whitehaven the workings extend a mile under the ocean, at about 600 feet below its bottom ; and it is asserted that the quality of the coals is still improving as the miner advances in this direction.

From this great deposit of coal alone, it is calculated that 28 millions of tons are raised annually ; nevertheless, it is also calculated that enough yet remains for the consumption of 1000 years to come.

A mere list of the strata that have been cut through to the depth of the mines at Whitehaven and Newcastle, would form a dry and uninteresting detail.

In the Restoration pit in St. Antony's colliery at Newcastle, which is 810 feet deep, the miner passed through 73 strata of various substances; of which 16 are coal. The first 6 strata of coal do not exceed 8 inches in thickness; 2 are 1 foot thick; 6 varied from 1 foot 6 inches to 4 feet; 1 is of 6 feet; and the thickest, which is the lowest, is of the thickness of 6 feet 6 inches.

In a mine at Preston Hows at Whitehaven, which is 642 feet deep, the miner passed through 117 strata, of which 17 are of coal; the thickest of these, which is the lowest, is 7 feet 10 inches.

All that can be said upon the origin of coal must be theoretical, and perhaps very remote from the truth: it is, however, certain, that there are few, if any, varieties of coal, which do not present more or less of the texture of wood. This appearance may be traced from the bitumenized wood, which still bears, though approaching in its nature to coal, the trunk, the branches, and even in some instances the very leaves of trees, through all the varieties of coal, into the most compact, slaty kind, of the oldest formation. In some, particularly from certain districts, the fibrous texture of wood is certainly remarkable; and the greater part of those who have given their attention to the probable origin of coal,

consider it to have resulted from vegetable remains.

That there still exist great underground forests is unquestionable. That in Prussia, yielding amber, may be quoted as an instance: and if coal be really of vegetable origin, this forest may hereafter in part yield important service to man, in the shape of coal. The immense bridge, not less than three miles in length, now existing and still receiving additions, on the Missouri, a river in North America, and consisting wholly of the trunks of trees, stopped in their progress down that river, may furnish coal to nations after the lapse of thousands of years.

But though some varieties of coal seem unquestionably of vegetable origin, certain others are not referable to it so decidedly: especially the slate coal of the oldest formation. This is frequently found in fragments of determinate shapes, particularly in that of a four-sided rhomboidal prism, presenting angles agreeing perfectly in measurement with those of crystals of mica.

Upon the whole, it must be acknowledged that the origin of coal is yet but little understood.



OF VOLCANOES.

On the subject of volcanoes, so much both of fact and of theory has been written, that it is

extremely difficult to compress into a narrow compass, any satisfactory detail respecting them or their origin.

Scientific writers have divided volcanoes into two kinds : pseudo or false volcanoes ; and true volcanoes.

Pseudo volcanoes usually occur in low situations, sometimes in hilly country. They are discoverable by a sensible heat, sometimes by smoke ; more seldom by flame. Sulphureous deposits and warm springs occur in their vicinity.

Pseudo volcanoes are almost always situated in the independent coal formation, and are considered to be caused by the spontaneous or accidental inflammation of beds of coal. Volcanoes of this description are to be observed in Bohemia, in Scotland, in England, and many in Kamschatka.

Of *true volcanoes* the number is considerably great. From the accounts of travellers, it appears that the whole number now in existence, most of which are occasionally in a state of activity, amounts to 193. They are thus distributed, according to Jameson, but it seems doubtful whether some pseudo volcanoes are not included.

Continent of Europe	1
European islands	12
Continent of Asia	8
Asiatic islands	58
African islands	8
Continent of America	87
American islands	19

In almost every country hills are found, which, from their shape, and the character of the surrounding masses, have, by some authors, been considered as extinct volcanoes: many of these are no longer supposed to be of volcanic origin; but it is certain that in Auverge, and in some other districts in France, as well as some in Spain, the remains of volcanoes are still to be seen; because the nature of their masses is indisputably volcanic.

Volcanoes are found at almost every elevation between the level of the sea and that of Cotopaxi in South America, which is 18,880 feet above it. Indeed, in many instances, volcanoes have burst from the bottom of the sea. Not long since, a considerable island was thus formed in one night, in the great Southern ocean; and many islands in the Archipelago, as, for instance, that called Santoniri, which is eight miles long, owe their origin to submarine volcanoes. The island of Teneriffe is about 45 miles in length, and 20 in breadth. There is an interesting and valuable memoir in the 2d vol. of the Geological Transactions,* describing a visit to the summit of its peak, which rises to an elevation of 11,000 feet above the sea. It appears, that in one district of this island there are 7 cones, exhibiting no traces of culture, no appearance of vegetation; that the soil of the island is altogether volcanic; that in one valley there are 100 strata of lava; and that every rock and stratum—in a word, the whole island, is the production of volcanic eruptions.

* By the Hon. Henry Grey Bennet, M. P. Pres. Geological Soc.

But our present object is with volcanoes, as being the cause of the ruin, and as forming anew, some of the constituent masses of the crust of the globe. Therefore, after some general and concise account of the nature of true volcanoes, we shall proceed to the consideration of the geological nature of one or two mountains, wherein those are situated of which we have the most authentic accounts, as of Vesuvius and *Ætna*; concluding our slight outline, (for it is a slight outline alone that I shall be able to present,) with such an account of the probable origin of volcanoes as the researches of the scientific afford.

Volcanoes have usually a conical shape, and are provided sometimes with one, sometimes with several mouths or craters.

In an active state, they occasionally eject smoke, vapour, flame, glowing and melted masses, and more rarely, water.

The occasional eruptions of large volcanoes are usually accompanied with earthquakes and lightning.

But when a volcano ejects only smoke, it is considered to be in a state of rest. This smoke is said to be composed of steam, or watery vapour, the muriatic and sulphureous gases; also of azote, carbonic acid, and hydrogen gas. If the smoke be black, it contains much carbonaceous matter; when gray, or of a white colour, it is principally composed of aqueous vapour.

In a greater state of activity, glowing masses

are ejected. These do not follow at regular periods, except in one instance that will be noticed presently. They are generally accompanied by a noise, proportioned in loudness to the magnitude of the stones that are projected, and the height to which they ascend. Ashes also are occasionally thrown out in prodigious quantities.

But the most striking phenomena exhibited by volcanoes, are the flowing streams of melted matter called lava, from their craters. These are usually preceded by earthquakes.

The instance just now mentioned, as an exception to the rule that the eruptions of volcanoes are not periodical, is that of Stromboli, one of the Lipari islands.

According to Dolomieu, the crater of Stromboli does not exceed fifty paces in diameter. This volcano is mentioned by Pliny; and, it is said, that from time immemorial, its eruptions have taken place about every 7 or 8 minutes. 'I saw it dart,' says Dolomieu, 'during the night, at regular intervals of 7 or 8 minutes, ignited stones, which rose to the height of more than 100 feet, forming radii a little diverging; but of which the greater part fell into the crater, while others rolled even to the sea. Each explosion was accompanied by a burst of red flame. The stones ejected are of a lively red, and sparkle, having the effect of artificial fire works. The approach of the eruption is not announced by any noise or dull murmur in the interior of the mountain.'

Vesuvius is a mountain of about 30 miles in circumference, and 3600 feet in height. The first recorded eruption is that of the year 79, which covered the towns of Pompeii and Herculaneum with a shower chiefly of sand and ashes of 80 feet in depth: but as late researches beneath this bed of ashes, and even to the foundations of the houses of Pompeii and Herculaneum, have proved that their streets were paved with volcanic matter, no doubt can be entertained that Vesuvius was a volcano at a much earlier period. Since the eruption just noticed as having happened in the year 79, thirty different eruptions of Vesuvius, of different degrees of violence, have been recorded.

In 1538, a mountain, principally of sand and ashes, 3 miles in circumference, and $\frac{1}{4}$ of a mile in height, was thrown up in one night.

An eruption of great violence, but inferior in this respect to that just noticed, was particularly described by Sir William Hamilton. It occurred in 1767, and lasted, at intervals, several days and nights. Streams of lava flowed, and prodigious quantities of ashes ascended, which, it is said, fell in Manfredonia, 100 miles distant from Vesuvius, in two hours after they were projected. Vast masses of stone were likewise thrown out by this eruption, which was accompanied by earthquake and lightning; many of these masses were measured by Sir William Hamilton; the largest of them was 108 feet in circumference, and 17 in length, but there were several not much inferior in bulk.

A stream of lava issued after the eruption of 1784, one mile wide and about twelve feet deep.

A particular account of volcanic eruptions would prove highly interesting; but our business with volcanoes is geological. I shall, therefore, content myself with saying, in respect to the nature of the eruptions of *Ætna*, that the projected matters much resemble those of *Vesuvius*. *Ætna* is above 10,000 feet high, and is about 130 miles in circumference: it was noticed as a volcano by *Diodorus Siculus*, 450 years before the Christian era.

But there is so great a difference in the nature of the substances thrown out by some of the American volcanoes, that they merit particular notice. Several of these are situated in mountains more than 10,000 feet high, and that of *Cotopaxi*, which is the highest, is 18,880 feet above the sea. According to *Humboldt*, these volcanoes scarcely ever throw out lava, but chiefly slag, ashes, a substance resembling pumice, and vast quantities of water, with an earthy or slimy mass, which often contains vast numbers of fishes. Hence, in the accounts of the tremendous volcanic eruptions that have taken place in the province of *Quito*, we hear only of overflowings, or of bodies being buried in slimy mud; never of the burnings that characterize the eruptions of some other volcanoes. When the volcano of *Carguairazo* fell down, on the night of the 16th July, 1698, it overflowed a tract of country 16 or 18 square miles,

with slimy mud. The number of human beings destroyed was so great, that the bodies were interred in heaps. During the great earthquake of the 4th February, 1797, 40,000 human beings were destroyed by the water and mud that issued from the mountains.

The substances ejected by pseudo volcanoes are, burnt clay, porcelain-jasper, earth-slag, columnar clay, iron-stone, and polishing slate.

Those ejected by true volcanoes are, granular limestone imbedding various minerals, most of them crystallized, and occasionally granite, mica-slate, greenstone, and sandstone: these substances have not undergone fusion, and frequently contain crystals of various substances unaltered; pumice, obsidian or volcanic glass, slime called volcanic tuff, sulphur, and muriate of ammonia, are, in general, in greater or less quantity, also the products of volcanic eruptions.

Of the origin of volcanoes, we can only reason from the nature of the substances they present to us. Many have been of opinion that volcanic mountains are wholly the results of the matter and masses ejected during eruptions, and that the real seat of volcanoes is very deep in the earth. Of the depth at which they may be situated, we know but little: but there seems no reason for believing that volcanic mountains, in the general, have been thrown up by the volcanoes that yet continue within them, or beneath their base.

Of the actual nature of Vesuvius as a mountain,

we know but little : we know that it often ejects calcareous matter, and that the country all around it is calcareous : but our information in regard to *Ætna* is more to the point. It has already been mentioned, that on the sides of that mountain a bed of sea shells has been discovered 2400 feet above the sea. It is therefore incredible that its whole bulk, or even that its whole surface, should be of volcanic origin, since the ruins of a temple built before our era, still stand on its side, uncovered by volcanic matter : and from the nature of the masses composing the volcanic mountain which fell in America, geologists are perfectly assured that the mountain itself could not have resulted from volcanic eruptions.

It has already been noticed, that no known volcano is seated in granite, and that granite is not to be seen near any volcano, except in very low situations. The same may be said of primitive rocks in general. One eruption of *Ætna* covered a space of 50 leagues in circumference, and 12 feet in thickness, with calcareous sand ; and calcareous earth, though it forms a very large proportion of secondary, enters very sparingly into the composition of primitive rocks : so that some geologists are not inclined to believe that volcanoes are very deeply seated in the earth.

Where fire exists we naturally believe the existence of combustibles. The known combustion of sulphur and iron filings, induced many to suppose that volcanoes originate from the decompo-

sition of vast deposits of pyrites, which are composed of iron and sulphur : but no spontaneous inflammation of pyrites, has ever been observed.

Werner has given his attention to the subject of the probable origin of volcanoes ; and knowing that coal has ignited spontaneously, and that there are vast beds of this substance, of which that at Liege in France, 90 feet thick, and 2 beds in Bohemia, one of which has been worked 90 feet deep, and the other much deeper, without reaching the bottom, are instances ; he is inclined to suppose, that to vast deposits of coal may be attributed the origin of volcanoes. The consequence of the spontaneous combustion of immense beds of coal, would be the melting of the stony beds that rest upon it. But the question arises, whence the power of raising vast masses, and of ejecting them, together with prodigious showers of sand ? The answer follows : the expansion of aqueous vapour, or steam ; and that water plentifully flows from many volcanoes, particularly those of America, we have already quoted sufficient evidence.

Humboldt, who, it must be acknowledged, has had ample opportunity for observing the phenomena of volcanoes, seems to be of opinion that certain of them are very deeply seated in the earth. He says it is a remarkable fact, that the volcanoes of Mexico are ranged in a line from east to west, forming a parallel of great elevation. In reflecting on this fact, and comparing it with

his observations upon certain phenomena attending Vesuvius, he is tempted to suppose that the subterraneous fire has pierced through an enormous crevice, which exists in the bowels of the earth, between the latitudes of $18^{\circ} . 59 .$ and $19^{\circ} . 12$, and stretches from the Pacific to the Atlantic ocean.

From the great differences in the theories of these naturalists, we are necessarily led to a conclusion, that as yet we have no sufficient data to enable us reasonably to account for the origin of volcanoes.



OF THE DELUGE.

THE traditions recorded by many persons and nations perfectly agree with the writings of Moses, and with the researches of geologists, in regard to the actual state of the surface of the globe, in this, that it has suffered a great catastrophe by water. It is alluded to by Berossus in the time of Alexander; by Plato; by the Hindus; who, according to Sir William Jones, mention it in one of their sacred books, or vedas, nearly in the same terms, and refer it to nearly the same period as Moses; and in the records of the Chinese philosopher Confucius, there is an allusion in the following terms: 'Having raised himself to heaven, Yao bathed the feet even of the highest mountains, covered the less elevated hills, and rendered the plains impassable:' allusions to the deluge are

also said to be involved in the astronomical calculations of the Chaldeans. It is remarked by Cuvier that *mere chance* could not give so striking a resemblance between the traditions of the Assyrians, the Hindus, and the Chinese; who moreover attribute the origin of their respective monarchies to the same period of about 4000 years back. The ideas of these three nations, who have so few features of resemblance, or rather, who are so entirely dissimilar in language, religion, and laws, could not have so exactly agreed on this point, if it had not been founded in truth. Other and remarkable traditional evidence may be adduced.

Acosta, in his history of the Indies, says that the Mexicans make particular mention of a deluge by which all men were drowned; and Dr. Watson, in one of his sermons, records a reply given by an inhabitant of Otabeite to one of our circumnavigators, to a question regarding their origin: that 'a long time ago, the earth was dragged through the sea; and their island, being broken off, was preserved.'

If it be true, as stated by Don Ulloa, that, during his travels in Peru, he found shells on a mountain more than 14,000 feet above the level of the sea, it affords a presumptive evidence that the sea must have attained that elevation; which is nearly equal to two-thirds the height of the most elevated points of land: and if it arose to so great a height, there seems no reason for assuming that as the

limit; we may reasonably thence infer that it covered the tops of the highest mountains.

There are, however, many circumstances relating to the lower parts of the earth, that afford interesting evidence of the great catastrophe. The compact argillaceous substance called shale, which lies over the independent coal formation, is often found to enclose vegetable remains; and it is said to exhibit, near Coalbrookdale, the impressions of gigantic ferns and reeds peculiar to the American continent. Deposits of marine plants and animals have been found in the interior of the European and Asiatic continents, frequently in considerable abundance. In Corsica are found the remains of an animal exclusively belonging to Siberia. In the year 1771 the carcass of a rhinoceros, an inhabitant of the torrid zone, was found on the banks of the Vilhovi. The bones of the mammoth, which is a species of the elephant, are found in three of the four quarters of the globe; and those both of the African and Asiatic elephant, have been found in many places in England: they occur likewise in Italy, France, Germany, and Sweden. It is worthy of remark, that, although in the two cited instances, the carcasses of the animals occurred nearly whole, entire skeletons are seldom found, but mostly separate bones, widely dispersed: and it is said that between the countries now inhabited by these animals, and some of those in which their bones have been discovered, there are chains of mountains exceeding 9000 feet in elevation.

All these circumstances seem to speak a language so perfectly intelligible that it cannot be misunderstood ; that since the creation of animals the world has suffered by an universal inundation ; that ‘all the fountains of the great deep were broken up, and the windows of heaven were opened.’

The manner in which this deluge was accomplished, is a problem that has long occupied the imaginations of philosophers and naturalists.

The question is this: since the tops of the highest mountains, which are about four miles above the present level of the sea, were covered with water; whence could be derived a sufficient quantity to surround, to the depth of four miles, a globe 25,000 miles in circumference.

It is evident that in order to accomplish this, the rain of forty days and nights was not sufficient; for the context says that ‘all the fountains of the great deep were broken up.’

What precisely is meant by the *great Deep*, and where it was situated, still continues to be problematical. Attempts at its explanation have caused the fabrication of at least fifty theories of the earth, not one of which is satisfactory.

Some of these theories were briefly recounted on a former evening, as being more or less connected with the creation of the world.

Burnet imagined that before the deluge, the earth was a mere crust, containing a vast abyss of waters; which issuing, deluged the earth, forming mountains of the broken crust.

Whiston supposed the deluge to have been occasioned by a comet. But a bare recital of hypotheses so gratuitous would scarcely be amusing; for in the general, there is so little agreement between them and the known phenomena of nature, as resulting from *those laws which in point of fact constitute nature*, that they are no longer deemed worthy of a place in a rational mind. Other theorists, however, bent upon a more reasonable philosophy, have looked into nature for a solution of this interesting question.

Kirwan has assumed that, in addition to rain of forty days and nights, a prodigious rising took place in the waters of the great Southern Ocean, which, he sets out with saying, is the greatest collection of waters on the face of the globe; that this ocean was the great deep mentioned by Moses; and he supposes that Noah resided on its borders, for otherwise he could not have seen that the great abyss was opened. He conceives that these waters were impelled northwards, with resistless impetuosity, against the continent which at that time probably united Asia with America; tearing up and sweeping away the whole of that immense tract, with the exception of those islands which still remain.

Here, then, is supposed a suspension of some of the laws of nature, or how could so vast a body of waters have arisen from its natural level, and have torn up and swept away a whole connecting continent, by what Kirwan terms its resistless

impetuosity. I do not propose to follow him through all the ramifications of his theory, but it may not be amiss to notice the apparent impossibility that the ark, or the little world within it, could have sustained the horrible shocks that must have been given to it by surges equal in power, and frequent enough in repetition, to cause the destruction of a continent.

Others have resorted to another theory. It is known that there is at all times a passage of electric matter into the earth, or from the earth into the atmosphere; and it is presumed that the electric fluid contained in the atmosphere, is the agent which suspends *therein* the water which rises from the earth in vapour.

Now they suppose that natural causes were so influenced for the sake of producing the deluge, as that the air being divested of the electric fluid, universal and amazing torrents of rain fell during forty days and nights. And it is further supposed that as the earth contained a double portion of electric fluid, which experiment has proved to possess the power of raising water upon and above the earth, that the waters contained in its fissures and caverns, and even whole oceans, were thereby raised so far above their natural elevations, as to cover the tops of the highest mountains.

Thus has the sagacity of man (according to the present extent of his knowledge) been brought into action, in the endeavour to account for this

wonderful phenomenon ; and we have seen that even those who have attempted to explain it, in some sort through the agency of natural causes, have been compelled to call in aid of their theories, the miraculous intervention of the Great Author of Nature.



OF THE INTERNAL STRUCTURE OF THE EARTH,
AND OF THE PROBABLE AGENT EMPLOYED IN
PRODUCING THAT STRUCTURE.

THE evidence produced on former evenings has satisfactorily proved, as I trust, that the researches of geologists have warranted them in making three grand features in the division of the rocks or masses which compose the crust of the globe.

PRIMITIVE ROCKS :

Consist only of crystalline deposits ;

Contain no organic remains ;

Are found below all other rocks ;

Rise from the base through the centre, and form the summits of lofty mountains.

TRANSITION AND FLUTZ ;

OR

SECONDARY ROCKS :

Consist partly of crystalline, partly of mechanical deposits ;

Contain organic remains of shells ; some of them not now found in the seas ;

Are never found under primitive rocks.

ALLUVIAL DEPOSITES :

Consist of mechanical deposites ;

Result from the ruin of rocks ;

*Contain the shells of fish now found in the ocean,
and the bones of large land animals ;*

Are found above all other rocks.

Geologists, almost universally, have agreed upon the nature and structure of the two latter descriptions of rocks, and even upon the manner in which they were deposited. For as almost all the rocks denominated secondary, and those called alluvial, contain the shells of fish which once lived in the sea ; all agree that these rocks must have been formed since those fish lived ; and that, generally speaking, they were deposited by the sea.

But in the estimation of some, primitive rocks, which are so termed chiefly because they contain no organic remains, do not owe their origin to the same cause. For though all coincide in the belief that the earth, to a certain depth, has once been in a fluid state, of which its spheroidal figure is considered to be a proof, yet opinions, as to the nature of that fluid, are most completely at variance ; inasmuch as one party considers it to have been caused by the agency of *fire*, and the other by that of *water*.

These two parties are termed *volcanists* and *neptunists* ; or more familiarly by geologists, *Huttonians* and *Wernerians* ; because the late Dr. Hutton of Edinburgh was the great champion for

the agency of fire, and because Werner, now residing in Germany, espouses the opposite theory.

Dr. Hutton seemed to consider that the great question we have to decide, is, *how our vast continents have been raised so high above the level of the sea.* Arguing from certain chemical facts, which it is not my province to explain, Dr. Hutton thought himself justified in supposing that the older rocks, as granite and the rest, were formed from a mass in a state of fusion by fire: that, owing to the intensity of this fire, and the expansion consequent to that intensity, in the central parts of the earth, all those rocks which are termed primitive, on which all others rest, and which form, as it were, the rough frame-work of the earth, *were raised up, forming continents and islands.* He moreover supposed, that, in cooling, these vast masses suffered contractions, producing rents and fissures in them, which afterwards were filled from the central fire, with metallic and other substances; constituting what are now known by the name of mineral veins.

Contemplating the causes of destruction perpetually operating, as volcanoes, exposure to rains and air, and sea-floods, Dr. Hutton also imagined that these, by the destruction they cause, and the quantity of ruin they carry from the land, are perpetually laying, in the unfathomable regions of the sea, the foundations of new continents; which, gradually rising, will in turn destroy those now existing.

Dr. Hutton does not suppose the earth, as it now is, to have been an original creation; but that it has resulted from the ruin of other and former worlds: of which he emphatically says, 'we can find no vestige of a beginning, no prospect of an end.'

Werner, on the contrary, does not seem to consider that the great question we have to decide is, how our vast continents have been raised so high above the sea.

He considers primitive rocks to be precisely in the situation in which they were first deposited; and as these rocks are all crystalline, or composed of vast aggregations of crystals, that the substances of which these crystals are composed, being held in solution in a vast body of water, greater in elevation than the summits of our highest mountains, followed the laws of affinity, and were deposited, forming those masses termed primitive rocks:

That these rocks, during their consolidation, split, forming rents and fissures; that these fissures were filled from the ocean with mineral bodies, and are what we now term mineral veins.

Such are the outlines of the two theories which now divide the opinions of geologists. Upon their comparative merits, it will not perhaps become me to say much. Yet, although no opportunity has hitherto occurred, or perhaps ever will occur, for my examination of geological facts on a large scale, I am unwilling to leave the subject without expressing some opinion.

It may be true that the chemical facts on which they rely who espouse the theory of the agency of fire, may not have been altogether controverted; and that these facts may apply, by possibility, to the formation of certain rocks: but arguing generally, from what is known of the nature of fire and of fuel, from what may be learned by the examination of hand specimens, and from the results obtained by the application of chemical agency to their component parts, I find it impossible not to give a preference to that theory which ascribes the formation of primitive rocks to the agency of water. It appears to me, that an opposition to the agency of fire may reasonably be supported by certain phenomena presented to us by the component substances of granite and gneiss, the two oldest of the primitive rocks, which are, for the most part, compounded of regular crystals of quartz, felspar, and mica, irregularly disposed in so far as regards each other.

If we could once arrive at any certainty that a single crystal of either of these substances had, in any case, been formed through the agency of either fire or water, it would seem to form reasonable ground for attributing the whole to the same agency. If, for instance, we could show that a crystal of quartz had been formed in water, it would go near to determine the question; because as the crystals of quartz, felspar, and mica, aggregately forming granite, are irregularly disposed with regard to each other; and as the crystals of

quartz sometimes enclose lesser crystals of felspar or of mica, it seems clear that the same agency must be referable for the production of the whole.

Now it has been already quoted that professor Seigling discovered some crystals of quartz that were deposited from an aqueous solution in which silex predominated, termed the *liquor of flints*, and which had accidentally been left eight years unobserved: these crystals were hard enough to give fire by the steel. It is asserted that Bergman also obtained crystals of the same substance from an aqueous solution containing silex.*

These facts may not be deemed conclusive of the question; but we have, as it seems to me, strong corroborative evidence in addition.

Crystals of quartz are by no means uncommon, in which are enclosed both *water* and *air*. I possess two or three such specimens, as well as another, enclosing, together with *water* and *air*, a minute portion of a dark substance that seems to be *bitumen*; which, turn the crystal any way, always presents itself in the air, and swimming on the water.

Now if, in the production of these crystals, fire had been the agent, would not the water have passed off in the form of vapour, and the bitumen have been consumed? It seems impossible to suppose these crystals to have been formed by any other agency than that of water.

* See Note, page 109: see also Nicholson's Journal, vol. i. page 217, of quarto edition.

Other facts are worth observing. If a crystal of quartz be exposed to a red heat, it loses its brilliancy, together with about three parts in the hundred of its weight, and is no longer transparent. This loss, it should seem, can be no other than the loss of water taken up during crystallization. The same holds, in part, with respect to felspar; which, though it never is transparent, loses from two to forty parts in the hundred of its weight, when exposed to the same heat.

Facts so pointed and important in regard to two of the three substances composing the oldest of the primitive rocks, seem to furnish strong evidence, that in the reducing into their present state the substances composing the great masses which form the crust of the globe, water was the agent.



WE have now completed the object we had in view, in dedicating five evenings to the investigation of simple and compound mineral bodies; namely, *an outline of the great masses composing the crust of the globe; of their arrangement; and of their component substances.*

In this view I have as much as possible studied brevity. It may be well imagined that there are many facts, considerations, and inquiries, of which nothing has been said; facts in regard to the relative natures and positions of compound mineral bodies and masses; considerations and inquiries into which those facts would necessarily lead.

Enough, however, I trust, has been said to show that geologists have truth for their object. That faculty of genius which consists in invention no longer presides; the theories which attributed the origin of the globe to a portion of the sun struck off by a comet, and fifty others equally absurd, which by their splendour once dazzled mankind, are gone by and neglected. Patient and profound investigation has taken their place; producing research nearly to the summits of our most elevated mountains, and to the greatest depths to which the miner can descend.

But since it is computed that all our researches cannot extend further into our earth, than, by comparison, the thickness of the paper which covers a globe of three feet in diameter, is to that globe; it will follow that we cannot possess any knowledge of what the central parts of the earth are actually constituted.

We have, however, the most indubitable evidence that the crust of the globe has been subject to revolutions, both partial and general. We are assured by the numerous facts that have been quoted, and by far more numerous which yet remain, that the sea must have changed both its *place* and its *height*. Proofs have been adduced that animal life has repeatedly and largely fallen the victim to these terrible events; there seems reason to conclude that some animals have been destroyed by sudden inundations; that others have been laid dry in consequence of the bottoms of the sea being suddenly elevated; and that these

calamities have caused great changes in the outer crust of the globe. It seems also clear, that since these first and greater commotions, those which followed, uniformly acted at a *less depth* and *less generally*. We have seen that the researches of geologists have ascertained that *of those animated beings, of which the remains are inclosed in those rocks which immediately rest upon primitive rocks, the races have become extinct ; that the newer rocks contain the remains of animals more nearly approaching to, but not absolutely of the same species as those inhabiting our present seas ; but that the newest contain only the remains of such animals as now exist in the seas, together with the bones of large land and amphibious animals.*

Every part of the globe distinctly bears the impress of these great and terrible events. The appearances of change and ruin are stamped on every feature. Change and ruin *by which not a particle of the creation has been lost, but which have been repeated, and are distinctly marked by the genera and species of the organic remains they enclose.*

Thus, those fossils and petrifications which heretofore were carefully collected as curiosities, now possess a value greater than as *mere curiosities*. They are to the globe what coins are to the history of its inhabitants ; they denote the period of revolution ; they ascertain at least comparative dates.

If the inquiry should arise, What benefit has resulted from ruin so extensive and so general ?

the answer is obvious; *soil and fertility*. If for a moment we imagine a world composed only of those rocks which we call primitive, which bear no marks of ruin, enclose no organic remains; we know from the nature of their component substances, that their exposure to the action of the elements during very many ages, would scarcely so separate and disintegrate them, as to produce a soil capable of any considerable vegetation; in other words, would fit the earth to receive and to maintain an extensive and almost universal population. A large and fertile part of England is composed of the ruin of rocks to a considerable depth; and this greatly obtains in all the most level and most fertile countries. Are we not then in degree justified in assuming, that this great ruin was designed to fit and prepare the earth for the support of the numerous animal tribes that inhabit it; most especially for MAN; who, doubtless, from his superior intellectual endowments, has emphatically been termed 'the Lord of the Creation.'

But our inquiries into utility need not stop here. All our researches have evinced such unquestionable proofs of design and contrivance, that it is impossible not to see them; and if we see them, it is or ought to be equally impossible not to ascribe them to the great Artificer of the universe. This, indeed, is the reasonable end and aim of all our inquiries, and all our philosophizing.

Without *mountains*, what in all probability would be the earth? A swamp or a sandy desert;

and the atmosphere a receptacle of noisome and pestilential exhalation. As conductors of the electric fluid, mountains contribute to the production of rain, which fertilizes the earth and purifies the atmosphere. They are the principal repositories of metallic ores. Their benefits, therefore, are great and extensive.

Hitherto the labours of the chemist have discovered 27 *metals*, 9 *earths*, 2 *fixed alkalies*, and the two bases of combustible bodies, *sulphur and carbon*: and these (although some of them may possess some characters in common) have each some peculiar to themselves, and are, therefore, termed *elementary substances*. It may be remarked that these, either in the simple or compound state, are found in quantity *admirably apportioned to their utility*; and, in the same proportion, with whatsoever they may be combined, they are generally most readily and easily freed from those substances with which they are compounded. Is it possible for one moment to doubt whether all this *exhibits design and contrivance for the benefit of Man?*

But further; is not design manifest in regard to the depositions of salt and of coal, so essential to man? Suppose these to have taken place between the earlier rocks, or in the masses of primitive mountains, or any where except where we find them; that is, just beneath the surface: they would in that case have been nearly lost to man. Can we appreciate their present benefits? Can

we doubt that there was design in placing them where we find them ?

But, in speaking of soil and fertility as the consequence of the decay and ruin every where to be observed, I omitted one important consideration ; namely, the decay of vegetables, which so essentially contributes to fertility, as principally composing what we call *mould*. But there is one exception to the general rule in regard to the decay of vegetable matter, not sufficiently noticed, as being an exception in the favour of Man ; but which demands a thought. The leaf, after it has fallen, enriches our soil ; the gardener esteems it an excellent manure ; but the tree that bore the leaf, when deprived of life, decays not until after a long period of time ; not until it has long been in the service of Man. Without wood, how should we be able to accomplish the many purposes to which it is turned ? Where could we find a substitute ?

This, I grant, is a digression ; but it is almost the only one I have allowed myself. I shall perhaps be excused, when it is considered, that it is adding another proof, in itself rather too obvious to attract much notice, to the many already adduced, *that design and contrivance are every where manifest, and in every thing intended for the advantage of Man.*

THE END.




Branch Storage

7 DAY USE

RETURN TO DESK FROM

RETURN EARTH SCIENCES LIBRARY
TO → 230 McCone Hall 642-2997
LOAN PERIOD 1 2 3
1 MONTH
4 5



-168

U.C. BERKELEY LIBRARIES



C034635783

